

FORM PTO-1390		U S DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 1588.GLE.PT
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/ES/US) CONCERNING A FILING UNDER 35 U.S.C 371			U S APPLICATION NO (If known, see 37 CFR 1.5) 09/786835
INTERNATIONAL APPLICATION NO PCT/EP99/06692	INTERNATIONAL FILING DATE 11 September 1999	PRIORITY DATE CLAIMED 10 September 1998	
TITLE OF INVENTION REGULATORY SEQUENCES FOR ROOT SPECIFIC OR ROOT ABUNDANT GENE EXPRESSION IN PLANTS			
APPLICANT(S) FOR DO/EO/US Günter Feix, Doerte Wulff			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: <ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)). <ol style="list-style-type: none"> <input checked="" type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 10 (35 U.S.C. 371(c)(3)). <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). <input type="checkbox"/> A translation of the annexes to the International Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). <p>Items 11. to 16. below concern document(s) or information included:</p> <ol style="list-style-type: none"> <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input type="checkbox"/> A change of power of attorney and/or address letter. <input checked="" type="checkbox"/> Other items or information: <ul style="list-style-type: none"> - Certificate of Mailing - Postcard 			

U.S. APPLICATION NO. (If known, see 37 CFR 1.52)		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER			
09/786835							
<p>17. <input type="checkbox"/> The following fees are submitted:</p> <p>BASIC NATIONAL FEE (37 CFR 1.492 (a)(1) - (5)):</p> <p>Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$970.00</p> <p>International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO. \$840.00</p> <p>International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$690.00</p> <p>International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4). \$670.00</p> <p>International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4). \$96.00</p> <p style="text-align: center;">ENTER APPROPRIATE BASIC FEE AMOUNT -</p> <p>Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).</p>				CALCULATIONS		PTO USE ONLY	
						840.00	
TOTAL OF ABOVE CALCULATIONS =		\$1190.00					
Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).		\$					
SUBTOTAL =		\$					
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).		\$					
TOTAL NATIONAL FEE =		\$					
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +		\$					
TOTAL FEES ENCLOSED =		\$1190.00					
		Amount to be refunded:		\$			
		charged:		\$			

09/786835

JC02 Rec'd PCT/PTO 09 MAR 2001

- a. ☒ A credit card authorization in the amount of \$1190.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. 50-0881 in the amount of \$_____ to cover the above fees.
A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge additional fees which may be required, or credit any overpayment to Deposit Account No. 50-0881. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO

Randall B. Bateman
MORRISS, BATEMAN, O'BRYANT & COMPAGNI
5882 South 900 East, Suite 300
Salt Lake City, Utah 84121
(801) 685-2302 phone
(801) 685-2303 facsimile

SIGNATURE

Randall B. Bateman

NAME

37,774

REGISTRATION NUMBER

CERTIFICATE OF DEPOSIT UNDER 37 C.F.R. § 1.10

I hereby certify that this correspondence is being deposited with the United States Postal Service as First Class Mail, postage prepaid, under 37 C.F.R. § 1.10 on the date indicated below and is addressed to Asst. Commissioner for Patents & Trademarks, Washington, D.C. 20231.

Randall B. Bateman
Randall B. Bateman

March 9, 2001
Date

EXPRESS MAIL NO. EL 593621462 US

13 Rec'd PCT/PTO 11 APR 2001
09/786835

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Günter Feix et al.
Serial Number: _____(National Stage of PCT/EP99/06692)
Filed: 10 September 1999
Group: _____
Examiner: _____
For: REGULATORY SEQUENCES FOR ROOT SPECIFIC OR
 REOOT ABUNDANT GENE EXPRESSION IN PLANTS
Attorney Docket: 1588.GLE.PT

Commissioner of Patents
and Trademarks
Washington, D. C. 20231

I hereby certify that this correspondence is being deposited with the United
States Postal Service via First Class Mail addressed to:
Commissioner of Patents and Trademarks,

Washington, D.C. 20231 on

April 3, 2001
(Date of Deposit)

Randall B. Baler

PRELIMINARY AMENDMENT

Dear Sir:

Prior to examination of the above-referenced application, please enter the following
amendment.

In The CLAIMS.

Please cancel claim 1 through 29 and replace therefor, new claims 30 through 68 as set
forth in the replacement sheets attached hereto.


A credit card authorization to cover the additional claim fees is also included.

09/786835, 061901

Applicant submits that the application is believed to be in condition for allowance. Should the Examiner believe that any adverse action is necessary, it is requested that he contact Randall B. Bateman at (801) 6685-2302 so that such matters may be resolved as expeditiously as possible.

Respectfully Submitted,

MORRISS, BATEMAN, O'BRYANT & COMPAGNI, PC

A handwritten signature in black ink, appearing to read "Randall B. Bateman". The signature is fluid and cursive, with the first name "Randall" being more legible than the last name "Bateman".

Randall B. Bateman
Reg. No. 37,774

5882 South 900 East, Suite 300
Salt Lake City, Utah 84121
Telephone (801) 685-2302
E-mail: bateman@mboclaw.com

09786835-061901

CLAIMS

What is claimed is:

30. (New) A nucleic acid sequence for use in cloning and expressing a root specific nucleic acid sequence in a plant, selected from the group consisting of: a nucleic acid sequence of SEQ ID No. 1; a nucleic acid sequence complementary to SEQ ID No. 1; a nucleic acid sequence having a degree of identity of more than 70% to SEQ ID No. 1 or to the sequence complementary to SEQ ID No. 1; alleles of the nucleic acid sequence of SEQ ID No. 1; alleles of the complementary sequence; alleles of a nucleic acid sequence having a degree of identity of more than 70% to SEQ ID No. 1; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to SEQ ID No. 1.

31.(New) A nucleic acid sequence for use in cloning and expressing a root specific nucleic acid sequence in a plant, selected from the group consisting of: a nucleic acid sequence of SEQ ID No. 2; a nucleic acid sequence complementary to SEQ ID No. 2; a nucleic acid sequence having a degree of identity of more than 70% to SEQ ID No. 2 or to the sequence complementary to SEQ ID No. 2; alleles of the nucleic acid sequence of SEQ ID No. 2; alleles of the complementary sequence; alleles of a nucleic acid sequence having a degree of identity of more than 70% to SEQ ID No. 2; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to SEQ ID No. 2.

32. (New) A nucleic acid sequence for use in cloning and expressing a root specific nucleic acid sequence in a plant, selected from the group consisting of: SEQ ID Nos. 3-11; a nucleic acid sequence complementary to any of SEQ ID Nos. 3-11; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 3-11; a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 3-11; alleles of the nucleic acid sequence of any of SEQ ID Nos. 3-11; alleles of a sequence complementary to any of SEQ ID Nos. 3-11; alleles of a nucleic acid sequence having a degree of

09786835 061901

identity of more than 70% to any of SEQ ID Nos. 3-11; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 3-11.

33. (New) A nucleic acid sequence for use in cloning and expressing a root specific nucleic acid sequence in a plant, selected from the group consisting of: SEQ ID Nos. 12-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 12-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 12-15; a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 12-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 12-15; alleles of a sequence complementary to any of SEQ ID Nos. 12-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 12-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 12-15.

34. (New) A nucleic acid sequence derived from maize for use in cloning and expressing a root specific nucleic acid sequence in a plant, the sequence being selected from the group consisting of: a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

35. (New) A vector comprising a nucleic acid sequence selected from the group consisting of:

a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

36.(New) The vector of Claim 35, which is a bacterial or viral vector.

37.(New) The vector of Claim 35, wherein the nucleic acid sequence is operably linked to a gene of interest.

38. (New) The vector of Claim 37, wherein the gene of interest is a gene which confers a character of agronomic or industrial benefit.

39. (New) The vector of Claim 38, wherein the gene of interest is a gene for resistance to infection by a virus, a gene conferring resistance to an herbicide or insecticide, or a gene whose expression confers male sterility.

40.(New) The vector of Claim 37, which further includes regulatory elements directing or enhancing the expression of the gene of interest.

41.(New) The vector of Claim 40, wherein the regulatory elements are 5', 3', or 5' and 3' elements.

42. (New) The vector of Claim 41, wherein the 3' regulatory element is a poly A addition sequence of an NOS gene of *Agrobacterium tumefaciens*.

09786635-064904

43.(New) The vector of Claim 40, which further includes T-DNA.

44.(New) The vector of Claim 43, which includes the left border of T-DNA, the right border of T-DNA, or both the left and right borders of T-DNA.

45. (New) The vector of Claim 43, wherein the nucleic acid sequence is located within the T-DNA or adjacent to the T-DNA.

46.(New) The vector of Claim 45, wherein the nucleic acid sequence is in conjunction with the gene of interest and is located within the T-DNA or adjacent to it.

47.(New) A host cell containing a vector comprising a nucleic acid sequence selected from the group consisting of:
a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

48. (New) The host cell of Claim 47, wherein the nucleic acid sequence is operably linked to a gene of interest, and the gene of interest is a gene which confers a character of agronomic or industrial benefit.

49.(New) The host cell of Claim 48, wherein the gene of interest is a gene for resistance to infection by a virus, a gene conferring resistance to an herbicide or insecticide, or a gene whose expression confers male sterility.

09786835, 061901
T06T90, 5E898260

50.(New) The host cell of Claim 48, which further includes regulatory elements directing or enhancing the expression of the gene of interest.

51.(New) The host cell of Claim 50, wherein the regulatory elements are 5', 3', or 5' and 3' elements.

52.(New) The host cell of Claim 51, wherein the 3' regulatory element is a poly A addition sequence of an NOS gene of *Agrobacterium tumefaciens*.

53.(New) The host cell of Claim 50, which further includes T-DNA.

54.(New) The host cell of Claim 53, which includes the left border of T-DNA, the right border of T-DNA, or both the left and right borders of T-DNA.

55. (New) The host cell of Claim 54, wherein the nucleic acid sequence is located within the T-DNA or adjacent to the T-DNA.

56. (New) The host cell of Claim 53, wherein the nucleic acid sequence is in conjunction with the gene of interest and is located within the T-DNA or adjacent to it.

57.(New) The host cell of Claim 47, which is a plant, yeast, or bacterial cell.

58.(New) The host cell of Claim 57, which is a cell from a monocotyledonous or dicotyledonous plant.

59.(New) A cell culture including a host cell selected from the group consisting of plant, yeast and bacterial cells; said host cell containing a vector comprising a nucleic acid sequence selected from the group consisting of:
a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence

complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

60.(New) A plant including a host cell containing a vector comprising a nucleic acid sequence selected from the group consisting of:

a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

61.(New) A composition of matter comprising seeds, propagation material, harvest material, or plant tissue, wherein the composition includes a host cell containing a vector comprising a nucleic acid sequence selected from the group consisting of:

a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a

degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

62.(New) A composition of matter comprising seeds, propagation material, harvest material, or plant tissue derived from a plant having a host cell containing a vector comprising a nucleic acid sequence selected from the group consisting of: a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15.

63.(New) A method of genetically modifying a cell, comprising a step of transforming the cell with a nucleic acid sequence selected from the group consisting of: a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; and alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15; wherein a gene of interest contained in the vector is expressible in the cell.

64.(New) The method of Claim 63, wherein the cell is a plant, bacterial, or yeast cell.

65. (New) The method of Claim 63, containing a further step of regenerating the transformed cell to produce a differentiated plant.

66.(New) The method of Claim 63, wherein the step of transforming is accomplished by transfer of a vector or of nucleic acid sequences from a bacterium to the cell.

67. (New) The method of Claim 63, wherein the step of transforming of the cell further includes either a step of microinjection with a vector or nucleic acid sequence or a step of particle bombardment with a vector or nucleic acid sequence, such that the step of transforming is accomplished by direct intake of a vector or nucleic acid sequence.

68. (New) A method for isolating a root specific gene and/or a root specific regulatory element from a plant, in which nucleic acid sequences derived from the plant are screened using a nucleic acid sequence selected from the group consisting of : a nucleic acid sequence of any of SEQ ID Nos. 1-15; a nucleic acid sequence complementary to any of SEQ ID Nos. 1-15; a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15 or to a sequence complementary to any of SEQ ID Nos. 1-15; alleles of the nucleic acid sequence of any of SEQ ID Nos. 1-15; alleles of a sequence complementary to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to any of SEQ ID Nos. 1-15; alleles of a nucleic acid sequence having a degree of identity of more than 70% to a sequence complementary to any of SEQ ID Nos. 1-15; and a vector comprising any of the preceding nucleic acid sequences and alleles.



06-19-01

JCO3 Rec'd PCT/PTO

19 JUN 2001

09/786835

PCT
Box - SEQ

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Günter Feix et al.

Serial Number: 09/786,835 (National Stage of PCT/EP99/06692)

Filed: 10 September 1999

Group: _____

Examiner: _____

For: REGULATORY SEQUENCES FOR ROOT SPECIFIC
OR REOOT ABUNDANT GENE EXPRESSION IN
PLANTS

Attorney Docket: 1588.GLE.PT

Assistant Commissioner for Patents
Washington, D. C. 20231

I hereby certify that this correspondence is being deposited with the United States Postal Service via Express Mail addressed to:
Assistant Commissioner for Patents,

Washington, D. C. 20231 on

June 19, 2001
(Date of Deposit)

Sequence Listing under 37 CFR 1.82(g)

Statement under 37 CFR 1.821(f)

09786835-061901



A computer-readable copy of the Sequence Listing for the above-referenced case is hereby submitted under the provisions of 37 CFR 1.821-1.825.

Applicant hereby states that the information recorded in computer-readable form is identical to that of the paper copy filed with the application.

Applicant further states that the Sequence Listing contains no new subject matter.

Respectfully submitted this 17th day of June, 2001.

Morriss, Bateman, O'Bryant & Compagni

Randall B. Bateman
Reg. No. 37,774

106790-5898260

5/PRTS

09/786835

JC02 Rec'd PCT/PTO 09 MAR 2001

-1-

Description

The present invention relates to isolated regulatory elements, in particular promoters and 3' regulatory elements, that drive root specific or root abundant expression in plants, in particular in monocotyledonous plants, to vectors containing the regulatory elements, to host cells containing the vectors, to plants containing the host cells and to methods for isolating regulatory elements that contribute to tissue-preferred gene expression in plants.

Gene expression is considered to comprise a number of steps from the DNA to the final protein product. Initiation of transcription of a gene is generally believed to be the predominant controlling factor in determining expression of a gene. The transcriptional controls are generally located in relatively short sequence elements embedded in the 5'-flanking and/or 3'-flanking region of the transcribed gene with which DNA-binding proteins may interact. These DNA sequence elements serve to promote the formation of transcriptional complexes and eventually initiate gene expression processes. It is furthermore known that the regulation of gene expression often depends upon the development stage and the tissue specificity of the cell concerned. Thus, certain tissues of organisms such as plants may exhibit a metabolism and a protein composition different from other tissues of the plant or different from the same tissue in a different developmental stage.

09/786835 "05.1901

Controlling the expression of genes in transgenic plants is considered to have high commercial value. The ability to control gene expression is useful for conferring resistance and immunity to certain diseases or to modify the metabolism of the tissue. The transfer of heterologous genes or genes of interest into a plant under control of tissue-specific genes provides a powerful means of conferring selective advantages to plants and to increase their commercial value.

It is also considered particularly important to develop transcriptional regulatory units that direct gene expression selectively to root tissue. Root preferred gene expression will provide several advantages to a plant, such as alteration of the function of the root tissue; modification of the growth rate; resistance to root preferred pathogens, pests, herbicides or adverse weather conditions; as well as broadening the range of soils or environments in which said plant may grow. Root abundant or root specific gene expression would provide a mechanism according to which morphology and metabolism may be altered to improve the yield and to produce useful proteins in greater amounts.

Thus, there is a need to provide regulatory elements capable of directing transcription specifically in root tissue. WO97/44448 describes promoter elements conferring root preferred gene expression. The disclosed promoter elements were obtained from maize seedlings. However, the isolated promoter elements may not be suitable for any desired transformation problem.

Furthermore, WO94/02619 discloses regulatory sequences for root specific or root abundant gene expression derived from Brassica sp. However, this promoter was not shown to work in monocotyle-

09786835-061901

donous plants with an efficiency great enough to provide means for commercial use.

Thus, the technical problem underlying the present invention is to provide novel regulatory elements for use in cloning and expressing root specific or root abundant genes, in particular for use in monocotyledonous plants which provide a high expression efficiency together with a high tissue specificity.

The present invention solves the technical problem underlying the present invention by providing purified and isolated new nucleic acid sequences for use in cloning and expressing a root specific or root abundant gene in a plant which are selected from the group consisting of

- (a) the nucleic acid sequence set out in SEQ ID No. 1 or a part or complementary strand thereof,
- (b) a nucleic acid sequence which hybridizes to the nucleic acid sequence defined in SEQ ID No. 1 or a complementary strand thereof and
- (c) alleles or derivatives of the nucleic acid sequence defined in (a) or (b) or a complementary strand thereof.

In particular, the problem is solved by providing a purified nucleic acid sequence according to the above wherein the sequence is selected from the group consisting of (a1) the nucleic acid sequence set out in SEQ ID No. 2 or a part or complementary strand thereof, (b1) a nucleic acid sequence which hybridizes to the nucleic acid sequence defined in SEQ ID No. 2 and (c1) alleles or derivatives of the nucleic acid sequence defined in (a1) or (b1).

09786333-061901

In a further particularly preferred embodiment, the problem is solved by providing a purified nucleic acid sequence according to the above which is selected from the group consisting of (a2) the nucleic acid sequence set out in any one of SEQ ID Nos. 3 to 11, (b2) a nucleic acid sequence which hybridizes to any one of the nucleic acid sequences defined in any one of SEQ ID Nos. 3 to 11 and alleles or derivatives of the nucleic acid sequence defined in (a2) or (b2).

The invention also relates to purified nucleic acid sequences for use in cloning and expressing a root specific or root abundant gene in a plant that are selected from the group consisting of (a3) the nucleic acid sequence set out in any one of SEQ ID No. 12, 13, 14 or 15 or a part or complementary strand thereof, (b3) a nucleic acid sequence hybridizing to any one of the sequences defined in (a3) and (c3) alleles or derivatives of the nucleic acid sequences defined in (a3) or (b3).

The sequences set out in SEQ ID No. 1 to 11 comprise isolated 5' regulatory elements and are considered to be promoters or parts thereof, i.e. promoter elements. These sequences are capable of modulating, initiating and/or contributing to the transcription of nucleic acid sequences operably linked to them. In a preferred embodiment of the present invention these sequences may additionally contain at their 3' terminus the nucleotide sequence from position 1761 to 1780 as depicted in SEQ ID No. 14, which is CCTGGACTCG CTCACTGGCA.

The sequences set out in SEQ ID No. 12, 13 and 15 comprise isolated 3' regulatory elements and are considered to be 3' transcription regulatory elements, or parts thereof, for instance polyadenylation signals. The sequences of SEQ ID Nos. 12 and 15

09:36:35 "061901
100730"

linked in 5' to 3' direction to each other represent the complete 3' regulatory element of the present invention.

These sequences are capable of modulating, in particular contributing to, or terminating the transcription of nucleic acid sequences operably linked to them.

The sequence set out in SEQ ID No. 14 is a genomic clone comprising all of the above sequences except the 3' regulatory element of the SEQ ID No. 15.

The present invention also relates to nucleic acid sequences which hybridize, in particular under stringent conditions, to the sequences set out in any one of SEQ ID No. 1 to 15. In particular, these sequences have a degree of identity of 70% to the sequence of SEQ ID Nos. 1 to 15.

In the context of the present invention, nucleic acid sequences which hybridise to any one of the specifically disclosed sequences of SEQ. Id. Nos. 1 to 15 are sequences which have a degree of 60% to 70% sequence identity to the specifically disclosed sequence on nucleotide level. In an even more preferred embodiment of the present invention, sequences which are encompassed by the present invention are sequences which have a degree of identity of more than 70% or 80% and even more preferred more than 90%, 95% or 99% to the specifically disclosed sequences on nucleotide level.

The present invention also relates to nucleic acid sequences which encode proteins wherein the amino acid sequence of the proteins have a degree of identity of 66% to 90% on amino acid level, most preferably a degree of identity of more than 90%, 95% or 99% identity on amino acid level.

09/86835 DE 1901

Thus, the present invention relates to nucleic acid sequences, in particular DNA sequences which hybridise under the following conditions to the sequences specifically disclosed:

Hybridisation buffer: 1 M NaCl; 1% SDS; 10%
dextran sulphate; 100 μ g/ml ssDNA
Hybridisation temperature: 65° C
First wash: 2 x SSC; 0.5% SDS at room
temperature
Second wash: 0.2 x SSC; 0.5% SDS at 65°C.

More preferably, the hybridisation conditions are chosen as identified above, except that a hybridisation temperature and second wash temperature of 68° C, and even more preferred, a hybridisation temperature and second wash temperature of 70° C is applied.

Thus, the present invention also comprises functionally equivalent nucleic acid sequences to the sequences of any one of SEQ ID No. 1 to 15, in particular sequences which have at least homology to the sequence of SEQ ID No. 1 to 15. The invention also relates to alleles and derivatives of the sequences mentioned above which are defined as sequences being essentially similar to the above sequences but comprising, for instance, nucleotide exchanges, substitutions (also by unusual nucleotides), rearrangements, mutations, deletions, insertions, additions or nucleotide modifications and are functionally equivalent to the sequences set out in SEQ ID No. 1 to 15.

The nucleic acid sequences of the present invention are, in a preferred embodiment, derived from maize (*Zea mays*), most preferably from maize roots.

The nucleic acid sequences of the present invention are useful for cloning tissue specific, in particular root specific nucleic acid sequences, in particular regulatory elements and/or genes, in plants, in particular in monocotyledonous plants. Thus, the present invention provides the means for isolation of transcription regulatory elements that direct or contribute to tissue-preferred gene expression in plants, in particular in monocotyledonous plants, such as maize. The present invention also provides the means of isolating tissue specifically expressed genes, in particular root specifically expressed genes and their transcripts.

The nucleic acid sequences of the present invention are also useful for expressing genes of interest in plants, in particular in the roots of plants and especially in the roots of monocotyledonous plants such as maize or of dicotyledonous plants such as sugar beets (*Beta vulgaris*). Thus, the present invention provides the means to direct the expression of a gene of interest in a tissue-specific or tissue-abundant manner in roots, for instance in roots of seedlings. Accordingly, the proteins encoded by the gene of interest can be accumulated in roots. For instance, the promoter of the present invention is particularly useful in driving the transcription of heterologous structural genes that confer disease immunity or resistance to disease-susceptible roots.

In the context of the present invention, a number of terms shall be utilized as follows.

In the context of the present invention the term "disease" encompasses any adverse condition caused to a plant by a virus or

an organism other than itself, such as fungi, bacteria and insects.

The term "promoter" refers to a sequence of DNA, usually upstream (5') to the coding sequence of a structural gene, which controls the expression of the coding region by providing the recognition for RNA polymerase and/or other factors required for transcription to start at the correct site. Promoter sequences are necessary, but not always sufficient to drive the expression of the gene. A "promoter element" constitutes a fraction of the DNA sequence of the promoter.

A "3' regulatory element (or "3' end") refers to that portion of a gene comprising a DNA segment, excluding the 5' sequence which drives the initiation of transcription and the structural portion of the gene, that determine the correct termination site and contains a polyadenylation signal and any other regulatory signals capable of effecting messenger RNA (mRNA) processing or gene expression. The polyadenylation signal is usually characterized by effecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognised by the presence of homology to the canonical form 5'-AATAA-3', although variations are not uncommon.

"Nucleic acid" refers to a large molecule which can be single or double stranded, composed of monomers (nucleotides) containing a sugar, phosphate and either a purine or pyrimidine. The nucleic acid may be cDNA, genomic DNA, or RNA, for instance mRNA.

The term "nucleic acid sequence" refers to a natural or synthetic polymer of DNA or RNA which may be single or double stranded, alternatively containing synthetic, non-natural or al-

097866335 "061901

tered nucleotide bases capable of incorporation into DNA or RNA polymers.

The term "gene" refers to a DNA sequence that codes for a specific protein and regulatory elements controlling the expression of this DNA sequence.

The term "regulatory element" refers to a sequence located upstream (5'), within and/or downstream (3') to a coding sequence whose transcription and expression is controlled by the regulatory element, potentially in conjunction with the protein biosynthetic apparatus of the cell. "Regulation" or "regulate" refer to the modulation of the gene expression induced by DNA sequence elements located primarily, but not exclusively upstream from (5') the transcription start of the gene of interest. Regulation may result in an all or none response to a stimulation, or it may result in variations in the level of gene expression.

The term "coding sequence" refers to that portion of a gene encoding a protein, polypeptide, or a portion thereof, and excluding the regulatory sequences which drive the initiation or termination of transcription. The coding sequence or the regulatory element may be one normally found in the cell, in which case it is called "autologous", or it may be one not normally found in a cellular location, in which case it is termed a "heterologous gene" or "heterologous nucleic acid sequence". A heterologous gene may also be composed of autologous elements arranged in an order and/or orientation not normally found in the cell in which it is transferred. A heterologous gene may be derived in whole or in part from any source known to the art, including a bacterial or viral genome or episome, eukaryotic nuclear or plasmid DNA, cDNA or chemically synthesized DNA. The structural gene may constitute an uninterrupted coding region or

09786635 061901

it may include one or more introns bounded by appropriate splice junctions. The structural gene may be a composite of segments derived from different sources, naturally occurring or synthetic.

The term "vector" refers to a recombinant DNA construct which may be a plasmid, virus, or autonomously replicating sequence, phage or nucleotide sequence, linear or circular, of a single or double stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell, in particular a plant cell.

As used herein, "plant" refers to photosynthetic organisms, such as whole plants including algae, mosses, ferns and plant-derived tissues. "Plant derived tissues" refers to differentiated and undifferentiated tissues of a plant, including roots, shoots, shoot meristems, coleoptilar nodes, tassels, leaves, cotyledonous petals, pollen, ovules, tubers, seeds, kernels and various forms of cells in culture such as intact cells, protoplasts, embryos and callus tissue. Within the context of the present invention, the term "root" includes various groups of primary, lateral seminal or crown roots, and root tips. Plant-derived tissues may be in planta, or in organ, tissue or cell culture. A "monocotyledonous plant" refers to a plant whose seeds have only one cotyledon, or organ of the embryo that stores and absorbs food. A "dicotyledonous plant" refers to a plant whose seeds have two cotyledons.

As used herein, "transformation" refers to the process by which cells, tissues or plants acquire properties encoded on a nucleic

09786633 "061901

acid molecule that has been transferred to the cell, tissue or plant.

"Transformation" and "transferring" refers to methods to transfer DNA into cells including, but not limited to, particle bombardment, microinjection, permeabilizing the cell membrane with various physical (e.g., electroporation) or chemical (e.g., polyethylene glycol, PEG) treatments.

The term "host cell" refers to a cell which has been genetically modified by transfer of a heterologous or autologous nucleic acid sequence or its descendants still containing this sequence. These cells are also termed "transgenic cells". In the case of an autologous nucleic acid sequence being transferred, the sequence will be present in the host cell in a higher copy number than naturally occurring.

The term "operably linked" refers to the chemical fusion of two of more fragments of DNA in a proper orientation such that the fusion preserves or creates a proper reading frame, or makes possible the proper regulation of expression of the DNA sequences when transformed into plant tissue.

The term "expression" as used herein is intended to describe the transcription and/or coding of the sequence for the gene product. In the expression, a DNA chain coding for the sequence of gene product is first transcribed to a complementary RNA, which is often an mRNA, and then the thus transcribed mRNA is translated into the above mentioned gene product if the gene product is a protein. However, expression also includes the transcription of DNA inserted in antisense orientation to its regulatory elements. Expression, which is constitutive and possibly further enhanced by an externally controlled promoter fragment

03786633 "061901
106T90" 5E99260

thereby producing multiple copies of mRNA and large quantities of the selected gene product, may also include overproduction of a gene product.

The term "translation start codon" or "initiation codon" refers to a unit of three nucleotides (codon) in a nucleic acid sequence that specifies the initiation of protein synthesis.

The term "signal peptide" refers to the N-terminal extension of a polypeptide which is translated in conjunction with the polypeptide, forming a precursor protein and which is required for its entry into the secretory pathway. The signal peptide may be recognised by the mechanisms within the same species or unrelated species plants, necessary for direction of the peptide into the secretory pathway. The signal peptide may be active in roots, seeds, leaves, tubers and other tissues of the plant. The term "signal sequence" refers to a nucleotide sequence that encodes a signal peptide. The term "ER targeting signal" refers to the N-terminal extension of a polypeptide, which is translated in conjunction with the polypeptide forming a precursor protein and which is required for its entrance into the Endoplasmic Reticulum (ER) of a cell. The ER targeting signal may be recognised by the mechanisms within the same species or unrelated species plants, necessary for direction of the peptide into the ER and/or the vacuole of a cell. ER targeting signals may be active in roots, seeds, leaves, tubers and other tissues of the plant. The term "ER targeting sequence" refers to a nucleotide sequence that encodes the ER targeting signal.

A "tissue specific promoter" refers to a sequence of DNA that provides recognition signals for RNA polymerase and/or other factors required for transcription to begin, and/or for controlling expression of the coding sequence precisely within certain

097333 0401
100190 559250

tissues or within certain cells of that tissue. Expression in a tissue specific manner may be only in individual tissues, or cells within tissues, or in combinations of tissues. Examples may include tissue specific expression in roots only and no other tissues within the plant, or may be in leaves, petals, ovules and stamen, and no other tissues of the plant. Here, "tissue specific" is also meant to describe an expression in a particular tissue or cell according to which the expression takes place mainly, but not exclusively, in the tissue. Such an expression is also termed "tissue abundant".

"Selective expression" refers to expression mainly, preferably almost exclusively, in specific organs of the plant, including, but not limited to, roots, leaves, tubers or seed. The term may also refer to expression at specific developmental stages in an organ, such as in early or late embryogenesis or in seedlings. In addition, "selective expression" may refer to expression in specific subcellular locations within the cell, such as the cytosol or vacuole.

The term "root specific nucleic acid sequence" refers to nucleic acid sequences, i.e. genes, coding sequences and/or regulatory elements which are exclusively or mainly active in roots of plants, in particular those which direct or contribute to a root abundant or root selective expression of a protein. The term "root abundant nucleic acid sequence" refers to nucleic acid sequences, i.e. genes, coding sequences and/or regulatory elements which are mainly active in roots of plants, in particular those which direct or contribute to a root abundant or root selective expression of a protein.

09786635 061901

The present invention also relates to a vector comprising the nucleic acid sequences according to the above, in particular to a bacterial vector, such as a plasmid, or a virus.

In a particularly preferred embodiment of the present invention the nucleic acid sequences, i.e. the 5' and/or 3' regulatory elements of the present invention contained in the vector, are operably linked to a gene of interest which in this context may also be only its coding sequence, which may be a heterologous or autologous gene or coding sequence. Such a gene of interest may be a gene, in particular its coding sequence, conferring, for instance, disease resistance; drought resistance; insect resistance; herbicide resistance; immunity; an improved intake of nutrients, minerals or water from the soil; or a modified metabolism in the plant, particularly its roots. Such a modified metabolism may relate to a preferred accumulation of useful substances in roots, for instance sugars or, vice versa, in the depletion of substances undesirable in roots, for instance certain amino acids. Thus, in the context of the present invention, a gene of interest may confer resistance to infection by a virus, such as a gene encoding the capsid protein of the BWYV or the BNYVV virus, a gene conferring resistance to herbicides such as Basta®, or to an insecticide, a gene conferring resistance to the corn rootworm, a gene encoding the toxic crystal protein of *Bacillus thuringiensis* or a gene whose expression confers male sterility. A gene of interest includes also a coding sequence cloned in antisense orientation to the regulatory sequences directing its expression. Such a construct may be used specifically to repress the activity of undesirable genes in plant cells, in particular in roots, most preferably in primary, lateral seminal and/or crown roots. The gene of interest may also comprise signal sequences, in particular ER targeting sequences,

0976633106191

directing the encoded protein in the ER and eventually for instance in the cell wall, vascular tissue and /or the vacuole.

In a particularly preferred embodiment the vector defined above is comprised of further regulatory elements directing or enhancing expression of the gene of interest such as 5', 3' or 5' and 3' regulatory elements known in the art, for instance enhancers such as the 35S-enhancer. Regulatory elements considered in the present invention also encompass introns or parts of introns inserted within or outside the gene of interest.

In a particularly preferred embodiment of the present invention the 3' regulatory element is a transcription termination region, preferably a poly A addition or polyadenylation sequence, most preferably the poly A addition sequence of the NOS gene of *Agrobacterium tumefaciens*.

Thus, the nucleic acid sequences of the present invention are useful since they enable the root specific expression of genes of interest in plants, in particular monocotyledonous plants. Accordingly, plants are enabled to produce useful products in their roots. The nucleic acid sequences of the present invention may also be useful to regulate the expression of genes of interest depending upon the developmental stage of the transferred cell or tissue. Furthermore, the present invention allows the specific modification of the metabolism in roots.

In a particularly preferred embodiment of the present invention the vector furthermore contains T-DNA, in particular the left, the right or both T-DNA borders derived from *Agrobacterium tumefaciens*. Of course, sequences derived from *Agrobacterium rhizogenes* may also be used. The use of T-DNA sequences in the vector of the present invention enables the *Agrobacterium tumefaciens*

09786335-061901

faciens mediated transformation of cells. In a preferred embodiment of the present invention, the nucleic acid sequence of the present invention, together with the gene of interest and - optionally - further operably linked regulatory sequences, is inserted within the T-DNA or adjacent to it.

The present invention also relates to a non-biological, technical method of genetically modifying a cell by transforming it with a vector or a nucleic acid sequence according to the above, whereby the gene of interest operably linked to the nucleic acid sequence or sequences of the present invention is expressible in the cell. In particular, the cell being transformed by the method of the present invention is a plant, bacterial or yeast cell. In a particularly preferred embodiment of the present invention, the above method further comprises the regeneration of the transformed cell to a differentiated and, in a preferred embodiment, fertile plant.

The method to transform a cell with a vector according to the present invention may be any method known to effectively transfer nucleic acid sequences into a host cell, thereby allowing its expression therein and eventually the regeneration of the cell to a plant. In a particularly preferred embodiment of the present invention, the transfer of the nucleic acid sequence may be mediated by *Agrobacterium tumefaciens* or *rhizogenes*. However, any other method known to eventually achieve the same results may be used, such as direct uptake of nucleic acid sequences, microinjection of nucleic acid sequences, or particle bombardment.

The present invention also relates to host cells transformed with the vector of the present invention, in particular plant, yeast or bacterial cells, in particular monocotyledonous or di-

0976633-061901
T06T305933/50

cotyledonous plant cells. The present invention also relates to cell cultures, tissue, calluses, etc. comprising a cell according to the above, i.e. a transgenic cell.

Furthermore, the present invention relates to plants, harvest and propagation material, parts of plants, plant tissue, plant seeds, plant leaves, plant roots, leaves, or embryos containing a host cell of the present invention. These plants, plant tissues, plant seeds and plant parts are also called "transgenic". These plants or plant parts are characterised by, as a minimum, the presence of the transferred nucleic acid sequence of the present invention in their genome or, in cases where the nucleic acid sequences are autologous to the transferred host cell, are characterised by additional copies of the nucleic acid sequences of the present invention. Preferably, the nucleic acid sequences of the present invention are linked to genes of interest or antisense DNA normally not found in the transferred cell operably linked to the nucleic acid sequence of the present invention, at least not in that order or orientation.

Thus, the present invention also relates to roots, leaves, calluses, plant tissue, tassels, etc., nonbiologically transformed, which possess, stably integrated in the genome of their cells, a heterologous or autologous nucleic acid sequence containing a regulatory element of the present invention recognised by the polymerases of the cells of said plant parts and the gene of interest encoding a protein of interest or being an antisense construct.

Thus, the invention relates to plants, nonbiologically transformed, which possess, stably integrated in the genome of their cells, a heterologous or autologous nucleic acid sequence containing a regulatory element of the present invention recognised

09785835 064901

by the polymerases of the cells of said plants and a gene of interest encoding a protein of interest or being an antisense construct.

The teaching of the present invention is therefore applicable to any plant, plant genus, or plant species wherein the regulatory element of the present invention is recognised by the polymerases of the cells.

Finally, the present invention relates to a method for isolating or cloning a root specific gene and/or root specific regulatory elements, such as promoters whereby a nucleic acid sequence of the present invention is used to screen nucleic acid sequences derived from any source, such as genomic or cDNA libraries derived from plants, in particular monocotyledonous plants. The nucleic acid sequences of the present invention thereby provide a means of isolating related regulatory sequences of other plant species which confer root specificity to genes of interest operably linked to them.

Further preferred embodiments of the present invention are mentioned in the claims.

The invention may be more fully understood from the following detailed sequence descriptions which are part of the present teaching. The SEQ ID Nos. 1 to 15 are incorporated in the present invention. The positions indicated below refer to the sequence numbering of SEQ ID No. 14.

SEQ ID No. 1 is a partial DNA sequence of the zmGRP3 (glycine-rich protein: GRP) gene of Zea mays. The 1760 bp (base pair) sequence spans the region from and including position 1760 towards the 5' end, up to and including position 1.

09786635-061901

SEQ ID No. 2 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 1348 bp sequence spans the region from and including position 1760 towards the 5' end, up to and including position 413.

SEQ ID No. 3 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 1096 bp sequence spans the region from and including position 1760 towards the 5' end, up to and including position 665.

SEQ ID No. 4 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 746 bp sequence spans the region from and including position 1760 towards the 5' end, up to and including position 1015.

SEQ ID No. 5 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 554 bp sequence spans the region from and including position 1760 towards the 5' end, up to and including position 1207.

SEQ ID No. 6 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 406 bp sequence spans the region from and including position 1760 towards the 5' end, up to and including position 1355.

SEQ ID No. 7 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 190 bp sequence spans the region from and including position 1760 towards the 5' end, up to and including position 1571.

SEQ ID No. 8 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 1354 bp sequence spans the region from and including

09786633-061901
106799-3388260

SEQ ID No. 9 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 1497 bp sequence spans the region from and including position 1497 towards the 5' end, up to and including position 1.

SEQ ID No. 10 is a partial DNA sequence of the zmGRP3 gene of *Zea mays*. The 1550 bp sequence spans the region from and including position 1550 towards the 5' end, up to and including position 1.

SEQ ID No. 11 is a partial DNA sequence of the zmGRP3 gene of *Zea mays*. The 1710 bp sequence spans the region from and including position 1710 towards the 5' end, up to and including position 1.

SEQ ID No. 12 is a partial DNA sequence of the zmGRP3 gene of *Zea mays*. The 856 bp sequence spans the region from and excluding the translation stop codon at position 2558 - 2560 towards the 3' end up to position 3416.

SEQ ID No. 13 is a partial DNA sequence of the zmGRP3 gene of *Zea mays*. The 440 bp sequence spans the region from and excluding the translation stop codon towards the 3' end up to position 3000.

SEQ ID No. 14 is a large part of the genomic DNA sequence of the zmGRP3 gene of *Zea mays* comprising 3416 nucleotides and encompassing the coding region of the gene, a sequence of 20 bp between and excluding the translation initiation codon ATG (position 1781 - 1783) and the nucleotides CCA at position 1758

- 1760 and the regulatory elements identified in SEQ ID Nos. 1 to 13.

SEQ ID No. 15 is a partial DNA sequence of the zmGRP3 gene of Zea mays. The 1074 bp sequence spans the region from and excluding the most 3' located base "C" at position 3416 of SEQ ID No. 14 to the sequence motif "TCC" at position 4488-4490. Thus, the complete genomic DNA sequence of zmGRP3 comprises 4490 bp.

The invention is further illustrated by way of example and the following drawings.

Figure 1 shows a restriction map of the genomic clone zmGRP3 described in SEQ ID No. 14.

Figure 2 shows restriction maps of 5' deletion constructs of the present promoter element corresponding to SEQ ID Nos. 1 to 7.

Figure 3 shows restriction maps of 3' deletion constructs of the present promoter element corresponding to SEQ ID Nos. 8 to 11.

Figure 4 shows a northern blot analysis for various tissues of Zea mays.

Figure 5 depicts DNA constructs used for transient expression experiments.

Example 1: Cloning of the genomic zmGRP3 sequence

The genomic clone zmGRP3 was isolated by differential screening of a cDNA library made from maize coleoptile nodes, subsequent

09786933-061901
T06T00"EE98/60

isolation of a specific root specific cDNA and isolation of the corresponding genomic clone.

A cDNA library from maize coleoptilar nodes with emerging developing young crown roots was made by choosing 7-9 days old wild-type maize seedlings. mRNA from 30-40 of the wild-type coleoptilar nodes (about 15 mg per piece of tissue) was isolated with oligo(dT)-cellulose using the "Quick prep mRNA purification kit" (Pharmacia, Freiburg). Double stranded cDNA was prepared using the "Time Safer"-cDNA synthesis kit of Pharmacia Freiburg. After elimination of cDNAs with a size less than 0.4 kb (kilobases) the cDNAs were ligated into the phage vector λ -ZAP-II (Stratagene) using the ligation kit "Gig-Pack-Gold" (Stratagene, Heidelberg). The phages were plated on E. coli strain XL 1B. The obtained library had a titre of 3.7×10^6 pfu/ml. The inserts had an average size of 1.4 kb. The phages of the cDNA library were plated out in low density (about 1000-2000 pfu per agar plate (12 x 12 cm)). Per plate two nitrocellulose filters as replicas were prepared. Both filters were then separately hybridized, each with a different probe.

As probes radioactively labelled first strand cDNAs of mRNA were used which a) were derived from the coleoptilar nodes of wild-type seedlings (with formation of crown roots) and b) derived from coleoptilar nodes of rt-cs mutant seedlings (without formation of crown roots) (Hetz, et al. Plant Cell, 1996. Pp. 10, 845-857). The first strand cDNA was labelled using radioactivity and, for further analysis, digoxigenin.

Radioactive Labelling of first strand cDNA:

Materials:

ca. 1-3 μ g mRNA

5x RT-Buffer

250 mM Tris/HCl pH 8.5

40 mM MgCl_2
150 mM DTT
dNTP-Mix dTTP, dGTP, dATP, ea 5 mM
dCTP 1.3 mM
Random primer (0.8 $\mu\text{g}/\mu\text{l}$)
RNase-Inhibitor (30 U/ μl)
32P-dCTP (10 $\mu\text{Ci}/\mu\text{l}$)
SuperScript Reverse Transcriptase (Gibco, 100 U/ μl)

For standard labelling 1-3 μg of mRNA were dissolved in 25.5 μl of H_2O , denaturated for 10 minutes at 65°C , then cooled in ice water. The following agents were added in order: 10 μl 5 x RT-Buffer, 1 μl of Rnasin, 2.5 μl of dNTP-Mix, 5 μl of random primer, 5 μl (=50 μCi) ^{32}P -dCTP, and 1 μl of SuperScript Reverse Transcriptase. The preparation was mixed thoroughly and incubated for 2 hours at 42°C . A final purification was done using the "Nucleotide Removal Kit" (Quiagen, Hilden).

Plaques giving a signal on the filter hybridized with the wild-type probe, but not giving a signal on the filter hybridized with rt-cs probe were excised and analysed. An in vivo excision was carried out and the cDNA insert was cloned into plasmid pBluescript. Altogether, 32 plaques were excised and analysed. For the further analysis, the isolated plasmid DNA clone was labelled with digoxigenin and hybridized in a northern blot analysis against 10 μg total RNA from wild-type and mutant coleoptilar nodes.

The labelling with digoxigenin of first strand-cDNA was done as follows:

Materials:

ca. 1-3 μg mRNA

5x RT-Buffer

250 mM Tris/HCl pH 8.5

09706633 064904

40 mM MgCl₂
150 mM DTT
dNTP-Mix dCTP, dGTP, dATP, ea 5 mM
dTTP 1.3 mM
RNase-Inhibitor (Fermentas, 30 U/ μ l)
SuperScript Reverse Transcriptase (Gibco, 100 U/ μ l)
Digoxigenin dUTP (Boehringer)
Oligo dT (0.6 μ g/ μ l)
EDTA solution (0.2 M)
LiCl solution (4 M)
EtOH (100%, 70%)

The mRNA was denaturated for 5 minutes in 23.4 μ l of H₂O at 65°C, then rapidly cooled in ice water. The following agents were added in order: 10 μ l 5x RT-Buffer, 1 μ l of RNase-Inhibitor (ca. 20 units), 5 μ l of oligo-dT (final concentration 0.06 μ g/ μ l) 3.6 μ l of DIG-dUTP (final concentration 0.13 mM), 5 μ l of dNTP-mix and 2 μ l of SuperScript Reverse Transcriptase. The preparation was then incubated for 2 hours at 42°C. The labelled first strand cDNA was precipitated by adding of 2 μ l of EDTA solution, 5 μ l of LiCl solution and 130 μ l of 100% EtOH. Following a washing with 70% EtOH, the cDNA resuspended in 50 μ l of H₂O and stored at -20°C.

The northern analysis showed that the clone isolated hybridized with wild-type total RNA, but not with the mutant total RNA. As the mutant tissue differed from the wild-type tissue in the absence of crown roots, the expression of the isolated clone could be related to the formation of crown roots in maize seedlings. According to northern blot analysis, the clone was expressed exclusively in the cells of roots of young maize seedlings, in particular in the tissue. The clone was used to isolate a full-length cDNA clone. The full-length cDNA clone (zmGPR3 cDNA) was

in turn used to isolate a genomic clone zmGRP3 which is partially described in SEQ ID No. 14 as follows:

The sequence given in SEQ ID No. 14 comprises the 5' regulatory elements of the present invention, the coding region of the maize zmGRP3 gene and the 3' regulatory elements of the present invention except the most distal part of the 3' region which, however, is given in SEQ ID No. 15. The coding region of zmGRP3 spans from the translation initiation codon at position 1781 to 1783 to the translation termination signal TGA at 2558 to 2560. The 5' regulatory element of the present invention comprises at position 1712 up to 1719 the putative TATA-box "tataaata".

The sequences depicted in SEQ ID Nos. 1 to 7 comprise the putative TATA-box and are particularly useful for the purposes described in the present invention.

The 3' regulatory element of the present invention, in particular the elements described in SEQ ID No. 12 and 13 comprise at position 2957 the polyadenylation signal. The 3' regulatory element located 3' to SEQ ID No. 12 and 13 and depicted in SEQ ID No. 15 may prove particularly useful as expression enhancing element.

The coding sequence of zmGRP3 codes for a glycin-rich protein, whose transcripts accumulate exclusively in roots of young maize seedlings following developmentally specific patterns. The protein exhibits a hydrophobic domain at the N-terminal region followed by repeated glycin-rich motifs. The hydrophobic domain at the N-terminal region is believed to be part of an ER-type signal peptide. The obtained genomic clone is the first nucleic acid sequence encoding a glycin-rich protein expressed exclusively in the root system, in particular that of maize.

00786635 "061901
T06T99" 5E999260

Example 2: Northern analysis

As the tissues to be compared, i.e. wild-type and mutant tissues differed only in the presence or absence of crown roots, the transcription of zmGRP3 mRNA could be identified to be closely connected with the formation of crown roots. However, to determine whether the zmGRP3 transcription could also be detected in other tissues or other types of roots, primary roots, lateral seminal roots, and lateral roots (of maize) the following northern analysis was carried out. mRNA was isolated from various organs such as kernels, shoot meristems, leaves, coleoptilar nodes, tassels, various zones of primary, lateral, seminal and crown roots and root tips of lateral roots. The mRNA was then on a gel and hybridized with the zmGRP3 cDNA isolated above.

Figure 4 A to D illustrate the detection of zmGRP3 transcripts in northern analysis. After electrophoretic separation and blotting, the RNAs (10 μ g) were hybridized with the DIG-labelled zmGRP3 cDNA. Total RNAs were isolated from; Fig. 4A, coleoptilar node with emerging young crown roots (lane 1), shoot meristem (lane 2), root tips of crown roots (lane 3), of primary roots (lane 4), or of lateral seminal roots (lane 5), leaves (lane 6), kernel (12-14 days after pollination, lane 7), or tassel (lane 8). Fig. 4B, C and D represent RNA samples taken from defined root segments of crown roots (B), of primary roots (C), or of lateral seminal roots (D): root tip up to cm 1 (lane 1), cm 2 (lane 2), cm 3 (lane 3), and cm 4 (lane 4). Lane 5 in C represents root tips (up to 1 cm) of secondary roots. The lower parts of A, B, C and D show the ethidium bromide stained gels displaying the 18S and 26S rRNA bands.

Figure 4B indicates that zmGRP3 mRNA accumulation is developmentally and spatially regulated within the root with the highest

level of accumulation occurring in the meristematic and elongation region (cm 1), low levels in early regions of cell maturation (cm2) and again increasing levels in later regions of cell maturation (cm 3 and 4) (cm sections as defined in Held et al, Plant Physiol. (1993) pp. 102, 1001-1008). In crown roots the zmGRP3 mRNA is predominantly expressed in the root tip. A weak hybridization signal is detected in the later stage of root development (cm 4). A possible explanation for this difference in accumulation between crown roots and primary/lateral seminal roots is that the development of primary and lateral seminal roots begins 4 to 5 days before crown roots. The transcript was surprisingly not detected in tips of lateral roots.

Example 3: Preparation of transgenic plants

Materials and Methods:

Isolation and culture of immature embryos for particle bombardment

Immature maize embryos (crosses between inbred line A188 and H99) with a size of 1.0-1.5 mm were isolated under sterile conditions. They were cultured with the scutellar side facing upward on modified N6-Medium (D'Halluin et al. 1993) with 1 mg/l 2,4-D, solidified with 0.8% agarose or 0.3% phytigel.

The explants were bombarded 4 to 12 days after isolation. For the osmotic pre-treatment, the embryos were placed with the scutellar side facing upward on MSC-medium (Zhong et al. 1992) in which a higher osmotic value was adjusted with sucrose to 670 mOsm/kg. The osmotic treatment was started 4 h before bombardment and was continued for 20 to 24 h after bombardment. In ex-

periments where the rhizobial part of the embryo was bombarded, the embryo was cut immediately or one day after isolation.

Particle bombardment

Particle acceleration was performed with the PDS 1000/He gun (BioRad, Munich, Germany). Five µg plasmid DNA was precipitated onto gold particles with an average size of 0.4-1.2 µm (Heraeus, Karlsruhe, Germany) following a protocol described by BioRad (Munich, Germany) and modified according to Becker et al. (1994). For stable transformation experiments, the particle-DNA pellet was resuspended in 240 µl ethanol. For each bombardment, 3.5 µl were spread on the macrocarrier.

Parameters used for particle bombardment

Distance between

A: rupture disk and macrocarrier	2.5 cm
B: macrocarrier and stopping screen	0.8 cm
C: stopping screen and target cells	5.5 cm
Gas pressure	1350 psi
Partial vacuum	28 inch Hg
Particles	gold (size: 0.4-1.2 µm)
Amount of particles	30 µg per shot

Constructs

In experiments for stable transformation, the p35Spat gene (Brettschneider et al., 1997) was used as selection marker gene. In co-transformation experiments 2.5 µg DNA of the plasmid with the selection marker gene was mixed with the second plasmid containing the promoter-reporter gene and precipitated together onto the particles. The promoter used for transient and stable

transformation experiments has the sequence depicted in SEQ ID No. 1 and was fused to a GUS reporter gene. Further deletion constructs in the promoter region were made which are depicted in Fig. 2 and 3.

Figure 2 depicts 5' deletion constructs (Fig. 2 (b)) of the full length promoter of SEQ ID No. 1 (Fig. 2(a)), while Figure 3 shows 3' deletion constructs (Fig. 3 (b)) of the full length promoter of SEQ ID No. 1 (Fig. 3 (a)).

In detail, Figure 2a is a schematic representation of the cloned promoter fragment of *zmGRP3*. The positions of the restriction sites of the following restriction enzymes are presented (the numbers denote the nucleotide positions of the restriction sites in the promoter fragment): X: *Xho*I (1), G: *Bgl* II (1206), V: *Pvu*I (120), K: *Kpn*I (1354), P: *Pst*I (588), N: *Bst*NI (572, 1760), S: *Sac*I (664), B: *Bam*HI (698, 1014), Z: *Bst*XI (412), C: *Cl*aI (1497, 1525, 1570), D: *Dra*I (906), Y: *Xba*I (963). Furthermore, the position of the putative TATA box is indicated.

Figure 2b gives an overview of the 5' promoter deletion clones prepared. Presented is in each case the length of the promoter fragment still present (boldface), as well as the size of the region deleted from the 5' end (in brackets) of the promoter. The fragments were cloned in a vector that contained the β -glucuronidase gene (*uid A*).

Figure 3a) is a schematic representation of the cloned promoter fragment of *zmGRP3*. The positions of the restriction sites of restriction enzymes are presented as in Fig. 2a.

Figure 3b) is an overview of the 3' promoter deletion clones prepared. Presented is in each case the length of the promoter

09363360
"06130"
09363360

fragment still present (boldface), as well as the size of the region deleted from the 3' end (in brackets) of the promoter. The fragments were cloned in a vector that contained the β -Glucuronidase Gen (uid A) downstream from the -46bp-region of the CaMV 35S promoter (with TATA box). Thus, these 3' deletion constructs contain the CaMV 35S promoter initiation site.

Expression studies

For transient expression studies, different maize tissues, i.e. roots, leaves of seedlings, germinating embryos, were bombarded with the promoter constructs identified above. For histochemical GUS-detection, the bombarded tissues were incubated one day after bombardment at 37°C for 20 h in X-gluc staining solution (McCabe et al. 1988). After staining leaf tissues with x-gluc chlorophyll was extracted in ethanol/acetic acid glacial (3:1) for 30 to 60 min at 65°C and washed once with 70% ethanol.

For quantification of the promoter strength, bombarded tissues were analysed in a protein GUS-assay (Jefferson et al. 1987). For standardisation of the transformation frequency, the promoter constructs were co-transformed with a luciferase reporter gene under control of the actin promoter from maize (positive control). As negative control, pUC 19 DNA was used in the same amount as the promoter constructs. One day after bombardment, proteins of the bombarded tissue were extracted and analysed for GUS and LUC activity. The analysed root tissue exhibited GUS and LUC activity, while the other bombarded tissues only showed LUC activity. These results clearly evidence root specificity of the promoter element of the present invention.

The table I below shows the results of the transformation of 5' deletion constructs.

09786635 "061901

Promoter	Root	Leaf	Embryo
SEQ ID No. 1, 0	+	-	-
SEQ ID No. 2, -412	+	-	-
SEQ ID No. 3, -664	+	-	-
SEQ ID No. 4, -1014	+	-	-
SEQ ID No. 5, -1206	+	-	-
SEQ ID No. 6, -1354	+	-	-
SEQ ID No. 7, -1570	+	-	-
negative control	-	-	-
positive control	+	+	+

Table I: Transformation of various maize tissues with 5' deletion promoter constructs.

All of the tested promoter constructs clearly and unambiguously show expression in roots, but not in leaves or in the embryo. Even the smallest promoter fragment (SEQ ID No 7, 190 bp length) shows significant expression exclusively in the root.

Figure 5 shows further gus-fusion constructs using the regulatory elements of the present invention (not to scale). The boxes marked "GUS" symbolise the coding region of the uidA gene from E. coli, the position of the start and stop codons is indicated in each case. The 35S promoter from pBI121 was deleted in the constructs produced (pBI121-35S) or replaced by the GRP3-promoter (pBI-GRP3-Prom, pBI121-GRP3-Prom-3') or the GRP3-promoter + 2x35S-Enhancer (pBI212-2x35S-En-GRP3-Prom-3'). In some cases moreover the 3' region of GRP3 (pBI121-GRP3-Prom-3', pBI121-2x35S-En-GRP-Prom-3') was inserted instead of the NOS-terminator. The positions of restriction endonuclease-sites relevant to this are indicated under the relevant constructs (H:

HindIII, L: Sal/I, X: XbaI, B: BamHI, M: SmaI, S: SacI, E: EcoRI).

Figure 5 thus shows further gus-fusions constructs using the regulatory elements of the present invention. PBI 121 is a positive control while PBI-121-35S is a negative control. PBI-121-GRP3-Prom is a construct using the promoter of SEQ ID No. 1 being functionally linked to a gus-reporter gene and the NOS-terminator. PBI-121-GRP3-Prom 3' is a DNA construct of the present invention using the promoter element of SEQ ID No. 1 functionally linked to the gus-reporter gene and a 3' regulatory element of the present invention having the sequence of SEQ ID No. 12 and added 3' thereto of SEQ ID No. 15. Therefore the 3' regulatory element spans the region from and excluding the translation stop codon at position 2558-2560 towards the 3' end up to position 4490. This 3' regulatory element being the combination of SEQ ID No. 12 and SEQ ID No. 15 has also been used in DNA construct PBI-121-2 X 35S-EN-GRP3-Prom-3' which is similar to the above described construct but additionally has the 2 X 35S-enhancer (nucleotide positions 7089 to 7357 and -346 to -78 of the CaMV virus genome relative to the start of transcription described in Fang et al. (1989) and Franck et al. (1980); data base accession No. V00141 Y 02048 VRL 12. Sept. 1993; NID: g 58821).

The following table II shows the expression results in transient essays in roots and leaves. Accordingly, it can be shown that both the 5' and 3' regulatory elements of the present invention contribute to tissue specific expression in roots indicating that DNA constructs using both the 5' and 3' elements of the present invention or the elements separately may prove useful in genetic engineering of plants.

Table II: Transient expression studies

Construct	Activity in the Transient Assay in Roots	Activity in the Transient Assay in Leaves
pBI121	+++++	+++++
pBI121-35S	-	-
pBI121-GRP3-Prom	+	-
pBI121-GRP3-Prom-3'	++	-
pBI121-2x35S-En-GRP3-Prom-3'	+++	-

(+: strong expression, degree indicated by number of "+", -: no or hardly any expression).

Explants were cultured after bombardment for two weeks on N6 1-100-25 medium at 26°C in the dark. Selection was started 14 days after bombardment when the explants were subcultured on N6 1-100-25 medium without casaminoacids supplemented with 5 mg/l phosphinotricin (PPT), the active compound of the commercial herbicide Basta®. After two weeks, the calli were subcultured for another two weeks on N6 1-100-25 medium with 5 mg/ l Basta®. For regeneration, embryogenic calli were transferred on hormonefree MS-medium with 1 mg/l PPT and cultured at 24°C under light (16 h).

Green shoots were transferred to Magenta boxes with half-strength MS-medium and 1 mg/l PPT solidified with 0.3 % phytagel. Plantlets with a height of 6-8 cm which survived selection were transferred into soil after they generated roots. Small plants were cultured in soil in the greenhouse for 1 week under a plastic or glass lid. Routinely, more than 90% of the regenerants survived the transfer into soil. To confirm the selection, surviving plants were sprayed once or twice with a Basta®-solution containing 250 mg/l PPT and 0.1% Tween.

09786635-061901

Transformants were verified by southern analysis

DNA isolation and Southern blot hybridisation

For the analysis of stable transformants, total genomic DNA was isolated from primary transformants and their progeny using the protocol of Dellaporta et al (1983). 10 or 15 µg of DNA, uncut or digested with restriction enzymes were separated by electrophoresis and transferred to a Hybond N membrane (Amersham/England) or a BiodyneA membrane (Pall/England). Introduced DNA of the present invention (SEQ ID No. 1) was detected using a modified protocol of the non-radioactive digoxigenin chemiluminescent method (Neuhaus-Url et al. 1993). Filters were hybridised with PCR-labelled Dig probes. Stably transformed transgenic maize plants could be obtained.

Example 4

In situ hybridization experiments of various types of maize roots with zmGRP

Method:

In situ hybridisation experiments were carried out according to a modified protocol from Coen et al., 1990 (Cell 63, 1311-1322). First the tissue to be analysed was fixed in 4% paraformaldehyde over night. The tissue then was dehydrated in an increasing concentration of ethanol and finally embedded in paraffin.

The tissue was cut in 8-12µm slices. After that the slices were hybridised with DIG labelled zmGRP3 RNA sense or antisense probes which had been hydrolytically digested to a size of about 200 bp. Hybridisation was carried out in 50% formamid at a tem-

09786835-061901

perature of 50°C. Washing and the immunological staining was carried out exactly as described by Coen et al., 1990.

Results:

Longitudinal sections of different 5 day old root types were used. These slices were derived from primary root as well as from lateral crown roots and from stem derived crown roots. The embryogenic primary root as well as lateral and stem derived crown roots showed the same hybridisation pattern. In all cases there was no signal with the sense probe, whereas with the antisense probe there was a strictly root specific hybridisation. This hybridisation was cell specifically and was restricted to the epidermal cell layer and cells of the columella root cap. In all other tissue types there was no hybridisation signal.

References

- Becker, D., Brettschneider, R. and Lörz, H. (1994) Fertile transgenic wheat from microprojectile bombardment of scutellar tissue. *The Plant Journal* 5, 299-307.
- Brettschneider et al (1997) Efficient transformation of scutella tissue of immature maize embryos. *Theoretical Applied Genetics* (94), 337.
- Coen et al., 1990 (*Cell* 63, 1311-1322).
- Dellaporta, S.L., Wood, J. and Hicks, J.B. (1983) A plant DNA minipreparation: Version II *Plant Mol. Biol. Rep.* 4, 419-21.
- D'Halluin, K., Bonne, E., Bossut, M., De Beuckeleer, M. and Lee-mans, J. (1993) Transgenic maize plants by tissue electroporation. *The Plant Cell* 4, 1495-1505.
- Fang et al., (1989) *Plant Cell* 1, 141-150
- Franck et al., (1989) *Cell* 21 (1), 285-294

McCabe, D.E., Swain, W.F., Martinelli, B.J. and Christou, P. (1988) Stable transformation of Soybean (Glycine max) by particle acceleration. Bio/Technology 6, 923-926.

McElroy, D., Zhang, W., Cao, J. and Wu, R. (1990) Isolation of an efficient Actin promoter for use in rice transformation. The Plant Cell 2, 163-171.

Neuhaus-Url, G. and Neuhaus, G. (1993) The use of the non-radioactive dioxigenin chemiluminescent technology for plant genomic Southern blot hybridisation: a comparison with radioactivity. Transgenic Research 2, 115-120.

Zhong, H., Srinivasan, C. and Sticklen, M.B. (1992) In-vitro morphogenesis of corn (Zea mays L.) Planta 187, 483-489.

0976635 06494
T06T90"SE99260

ART 34 AMDT

Gleiss & Große

Patentanwälte
München Stuttgart

31

Claims

1. A nucleic acid sequence for use in cloning and expressing a root specific nucleic acid sequence in a plant which is selected from the group consisting of
 - a) the nucleic acid sequence set out in SEQ ID No. 1 or a complementary strand thereof,
 - b) a nucleic acid sequence which has a degree of identity of more than 70 % to the nucleic acid sequence defined in (a) and
 - c) alleles of the nucleic acid sequence defined in (a) or (b).
2. The nucleic acid sequence of claim 1 which is selected from the group consisting of
 - a) the nucleic acid sequence set out in SEQ ID No. 2 or a complementary strand thereof,
 - b) a nucleic acid sequence which has a degree of identity of more than 70 % to the nucleic acid sequence defined in (a) and
 - c) alleles of the nucleic acid sequences defined in (a) or (b).

3. The nucleic acid sequence according to claim 1 or 2, which is selected from the group consisting of
- a) the nucleic acid sequence set out in any one of SEQ ID No. 3 to 11 or a complementary strand thereof,
 - b) a nucleic acid sequence which has a degree of identity of more than 70 % to the nucleic acid sequence defined in (a) and
 - c) alleles of the nucleic acid sequence defined in (a) or (b).
4. A nucleic acid sequence for use in cloning and expressing a root specific gene in a plant which is selected from the group consisting of
- a) the nucleic acid sequence set out in any one of SEQ ID No. 12 to 15 or a complementary strand thereof,
 - b) a nucleic acid sequence which has a degree of identity of more than 70 % to the nucleic acid sequences defined in (a) and
 - c) alleles of the nucleic acid sequence defined in (a) or (b).
5. The nucleic acid sequence of any one of the preceding claims, which is derived from maize.
6. A vector comprising the nucleic acid sequence of any one of the preceding claims.

31
-3-

7. The vector of claim 6, which is a bacterial or viral vector.
8. The vector of any one of claims 6 or 7, wherein the nucleic acid sequence of any one of claims 1 to 5 is operably linked to a gene of interest.
9. The vector of claim 8, wherein the gene of interest is chosen from a gene which confers a character of agronomic or industrial benefit.
10. The vector of claim 9, wherein the gene which confers a character of agronomic or industrial benefit is a gene for resistance to infection by a virus, a gene conferring resistance to a herbicide or to an insecticide or, alternatively, a gene whose expression confers male sterility.
11. The vector of any one of claims 6 to 10, which comprises further regulatory elements directing or enhancing the expression of the gene of interest.
12. The vector of any one of claims 6 to 11, wherein the further regulatory elements are 5', 3' or 5' and 3' elements.
13. The vector of claim 12, wherein the 3' regulatory element is the poly A addition sequence of the NOS gene of *Agrobacterium tumefaciens*.
14. The vector according to any one of claims 6 to 13, which furthermore contains T-DNA.

40
-A-

15. The vector of claim 14, which contains the left, the right or both T-DNA borders.
16. The vector according to claim 14 or 15, wherein the nucleic acid sequence is located within the T-DNA or adjacent to it.
17. The vector according to claim 16, wherein the nucleic acid sequence in conjunction with the gene of interest is located within the T-DNA or adjacent to it.
18. A host cell containing the vector of any one of claims 6 to 17.
19. The host cell of claim 18, which is a plant, yeast or bacterial cell.
20. The host cell of claim 19, which is a cell from a monocotyledonous or dicotyledonous plant.
21. A cell culture comprising a cell according to any one of claims 18 to 20.
22. A plant comprising a host cell according to any one of claims 18 to 20.
23. Seeds, propagation material, harvest material and plant tissue, comprising a host cell according to any one of claims 18 or 20 or derived from a plant according to claim 22.
24. A method of genetically modifying a cell by transforming a cell with a nucleic acid sequence according to any one of claims 1 to 5 or a vector according to any one of claims 6 to

T06F90" SE898/60

- 17, wherein the gene of interest contained in the vector is expressible in the cell.
25. The method of claim 24, wherein the cell is a plant, bacterial or yeast cell.
26. The method of claims 24 or 25, wherein the transformed cell is regenerated into a differentiated plant.
27. The method of any one of claims 24 to 26, wherein the cell is transformed by transfer of a vector or nucleic acid sequences from a bacterium to the cell.
28. The method of any one of claims 24 to 27, wherein the cell is transformed by direct intake of a nucleic acid sequence or a vector, by microinjection or particle bombardment with a nucleic acid sequence or a vector.
29. A method for isolating a root specific gene and/or root specific regulatory element from a plant, whereby a nucleic acid sequence of any one of claims 1 to 5 or a vector of any one of claims 6 to 17 is used to screen nucleic acid sequences derived from the plant.

Abstract

The present invention relates to regulatory elements for tissue specific, in particular root specific, expression of genes of interest in plants. The invention also relates to methods to obtain transgenic plants which show root specific expression of genes of interest and to methods for isolating root specific genes and/or root specific regulatory elements.

Fig.1




Fig.2

Fig.2a




Fig.2b





Fig.3

Fig.3a







Fig.5





23907

Declaration and Power of Attorney for Patent Application Erklärung für Patentanmeldungen mit Vollmacht

German Language Declaration

Als nachstehend benannter Erfinder erkläre ich hiermit an Eides Statt:

daß mein Wohnsitz, meine Postanschrift und meine Staatsangehörigkeit den im nachstehenden nach meinem Namen aufgeführten Angaben entsprechen, daß ich nach bestem Wissen der ursprüngliche, erste und alleinige Erfinder (falls nachstehend nur ein Name angegeben ist) oder ein ursprünglicher, erster und Miterfinder (falls nachstehend mehrere Namen aufgeführt sind) des Gegenstandes bin, für den dieser Antrag gestellt wird und für den ein Patent für die Erfindung mit folgendem Titel beantragt wird:

deren Beschreibung hier beigefügt ist, es sei denn (in diesem Falle Zutreffendes bitte ankreuzen), diese Erfindung

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Regulatory Sequences for Root Specific Or Root Abundant Gene

Expression In Plants

the specification of which is attached hereto unless the following box is checked:

☒ wurde angemeldet am 10 September 1999
unter der US-Anmeldenummer oder unter der
Internationalen Anmeldenummer im Rahmen des Vertrags
über die Zusammenarbeit auf dem Gebiet des Patentwesens
(PCT)
PCT/EP99/06692 und am
4 December 2000 abgeändert (falls
zutreffend)

☒ was filed on 10 September 1999
as United States Application Number or PCT International
Application Number
PCT/EP99/06692 and was amended on
4 December 2000 (if applicable).

Ich bestätige hiermit, daß ich den Inhalt der oben angegebenen Patentanmeldung, einschließlich der Ansprüche, die eventuell durch einen oben erwähnten Zusatzantrag abgeändert wurde, durchgesehen und verstanden habe.

Ich erkenne meine Pflicht zur Offenbarung jeglicher Informationen an, die zur Prüfung der Patentfähigkeit in Einklang mit Titel 37, Code of Federal Regulations, § 1.56 von Belang sind.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56.

German Language Declaration

Ich beanspruche hiermit ausländische Prioritätsvorteile gemäß Title 35, US-Code, § 119 (a)-(d), bzw. § 365 (b) aller unten aufgeführten Auslandsanmeldungen für Patente oder Erfinderurkunden, oder § 365(a) aller PCT internationalen Anmeldungen, welche wenigstens ein Land ausser den Vereinigten Staaten von Amerika benennen, und habe nachstehend durch ankreuzen sämtliche Auslands- anmeldungen für Patente bzw. Erfinderurkunden oder PCT internationale Anmeldungen angegeben, deren Anmeldetag dem der Anmeldung, für welche Priorität beansprucht wird, vorangeht.

Prior Foreign Applications
(Frühere ausländische Anmeldungen)

98117251.3 ✓	Europe ✓
(Number)	(Country)
(Number)	(Land)
_____	_____
(Number)	(Country)
(Number)	(Land)

Ich beanspruche hiermit Prioritätsvorteile unter Title 35, US-Code, § 119(e) aller US-Hilfsanmeldungen wie unten aufgezählt.

_____	_____
(Application No.)	(Filing Date)
(Aktenzeichen)	(Anmeldetag)
_____	_____
(Application No.)	(Filing Date)
(Aktenzeichen)	(Anmeldetag)

Ich beanspruche hiermit die mir unter Title 35, US-Code, § 120 zustehenden Vorteile aller unten aufgeführten US-Patentanmeldungen bzw. § 365(c) aller PCT internationalen Anmeldungen, welche die Vereinigten Staaten von Amerika benennen, und erkenne, insofern der Gegenstand eines jeden früheren Anspruchs dieser Patentanmeldung nicht in einer US-Patentanmeldung, bzw. PCT internationalen Anmeldung in in einer gemäß dem ersten Absatz von Title 35, US-Code, § 112 vorgeschriebenen Art und Weise offenbart wurde, meine Pflicht zur Offenbarung jeglicher Information an, die zur Prüfung der Patentfähigkeit in Einklang mit Title 37, Code of Federal Regulations, § 1.56 von Belang sind und die im Zeitraum zwischen dem Anmeldetag der früheren Patentanmeldung und dem nationalen oder im Rahmen des Vertrags über die Zusammenarbeit auf dem Gebiet des Patentwesen (PCT) gültigen internationalen Anmeldetags bekannt geworden sind

_____	_____
(Application No.)	(Filing Date)
(Aktenzeichen)	(Anmeldetag)

_____	_____
(Application No.)	(Filing Date)
(Aktenzeichen)	(Anmeldetag)

Ich erkläre hiermit, daß alle in der vorliegenden Erklärung von mir gemachten Angaben nach bestem Wissen und Gewissen der Wahrheit entsprechen, und ferner daß ich diese eidesstattliche Erklärung in Kenntnis dessen ablege, daß wissentlich und vorsätzlich falsche Angaben oder dergleichen gemäß § 1001, Title 18 des US-Code strafbar sind und mit Geldstrafe und/oder Gefängnis bestraft werden können und daß derartige wissentlich und vorsätzlich falsche Angaben die Rechtswirksamkeit der vorliegenden Patentanmeldung oder eines aufgrund deren erteilten Patentes gefährden können.

I hereby claim foreign priority under Title 35, United States Code, § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Priority Not Claimed
Priorität nicht beansprucht

September 11, 1998 ✓ ☐
(Day/Month/Year Filed)
(Tag/Monat/Jahr der Anmeldung)

_____ ☐
(Day/Month/Year Filed)
(Tag/Monat/Jahr der Anmeldung)

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and , insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

_____ ☐
(Status) (patented, pending, abandoned)
(Status) (patentiert, schwebend, aufgegeben)

_____ ☐
(Status) (patented, pending, abandoned)
(Status) (patentiert, schwebend, aufgegeben)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.



German Language Declaration

VERTRETUNGSVOLMACHT: Als benannter Erfinder beauftrage ich hiermit den (die) nachstehend aufgeführten Patentanwalt (Patentanwältin) und/oder Vertreter mit der Verfolgung der vorliegenden Patentanmeldung sowie mit der Abwicklung aller damit verbundenen Angelegenheiten vor dem US-Patent- und Markenamt: (Name(n) und Registrationsnummer(n) auflisten)

Julie K. Morriss 33,263 Randall B. Bateman, 37,774
David W. O'Bryant 39,793 Frank W. Compagni, 40,567

MORRIS, BATEMAN, O'BRYANT & COMPAGNI
5882 South 900 East, Suite 300
Salt Lake City, Utah 84121

Postanschrift:

Randall B. Bateman (801) 685-2302

Telefonische Auskünfte. (Name und Telefonnummer)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith: (list name and registration number)

Julie K. Morriss 33,263 Randall B. Bateman, 37,774
David W. O'Bryant 39,793 Frank W. Compagni, 40,567

MORRIS, BATEMAN, O'BRYANT & COMPAGNI
5882 South 900 East, Suite 300
Salt Lake City, Utah 84121

Send Correspondence to

Randall B. Bateman (801) 685-2302

Direct Telephone Calls to (name and telephone number)

Vor- und Zuname des einzigen oder ersten Erfinders <u>Günter Feix</u>	Full name of sole or first inventor <u>Günter Feix</u>
Unterschrift des Erfinders <u>Günter Feix</u> Datum <u>4.5.01</u>	Inventor's signature Date
Wohnsitz <u>Freiburg, Germany</u>	Residence <u>Freiburg, Germany</u>
Staatsangehörigkeit <u>German</u>	Citizenship <u>German</u>
Postanschrift <u>Rehagweg 13</u> <u>D-79100 Freiburg</u> <u>Germany</u>	Post Office Address <u>Rehagweg 13</u> <u>D-79100 Freiburg</u> <u>Germany</u>
Vor- und Zuname des zweiten Miterfinders (falls zutreffend) <u>Doerte Wulff</u>	Full name of second joint inventor, if any <u>Doerte Wulff</u>
Unterschrift des zweiten Erfinders <u>Doerte Wulff</u> Datum <u>4.5.01</u>	Second Inventor's signature Date
Wohnsitz <u>Freiburg, Germany</u>	Residence <u>Freiburg, Germany</u>
Staatsangehörigkeit <u>German</u>	Citizenship <u>German</u>
Postanschrift <u>Buchenstraße 10</u> <u>D-79102 Freiburg</u> <u>Germany</u>	Post Office Address <u>Buchenstraße 10</u> <u>D-79102 Freiburg</u> <u>Germany</u>

(In Falle dritter und weiterer Miterfinder sind die entsprechenden Informationen und Unterschriften hinzuzufügen)

(Supply similar information and signature for third and subsequent joint inventors)

-1-

SEQUENCE LISTING

<110> Agricultural Technology and Genetics GmbH

<120> Regulatory Sequences for Root Specific or Root Abundant
Gene Expression in Plants

<130> 23143 ATG

<140>

<141>

<150> EP 98117251.3

<151> 1998-09-01

<160> 15

<170> PatentIn Ver. 2.1

<210> 1

<211> 1760

<212> DNA

<213> Zea mays

<400> 1

```

ctcgagatgt tggatagaag aggaattgctg agcaaccgag gtctccaagg caccgccaacg 60
cgcacccctgc gcagttaggt ggtcgctgat gtgcggaagg agcttcgatt gttcgtcgat 120
cgcgagcttc agatttggct ccatgacccc agattgtttg gcgcggcgcg tgtaatgagc 180
agacgccaga cccgagaccg gtgtctccga taccaattgt tagcaatagc agcggacaag 240
cagcgaaaca tccgagagac atagtttggg ggaggagcag atagcttgga gaggaagaag 300
aacaatagt acagcgtaga ttcctctctg gtctcctctg ttaccctctc tcaatctaata 360
actcctatta caaggtcggc ccatcaggcg acccagtcgc ggttcacga acttgggatt 420
gggtttgtgc ggacgccgac tggttcttgg ctctgctgga gtcttgtctg tggagctgtt 480
gctcgcgctg ctaacaatct cttgttgttt gcgacgactt cgcgtggcca cgcgttaact 540
gcgctctgct caaagaaatt gcctatacag tgcctggcgt cagctgcagg cgttgaatcc 600
gaggtogcca gtacgaggca aacgctgagc tgcctgctgt accggttta ttaataactg 660
agctctactc gctacgttag ctataggcac cagaggcgga tccagagggg gggcaaagta 720
ggccatggcc cccctcaat tttgtacacc cttttataacc taatgattat ttagtattaa 780
gtgtagaaaa aatcaaaatt ttagatagta taaaaatctt gttctaaatc tctaaaaacta 840
atagctagtt gctaaaatta gccaaaagggt ttagaatgga tcaactaatt gttagctata 900
ccctttaaat aagtattaat tgtagatagg cccaacctag ctaaaaccag ttaacagtta 960
gctctagatg tttagaacat ggccctaaaca accatattat gacccactc gccggatcct 1020
ctcggttcag tcctagcatc taccagtctg actctaactc tcccgacacc acctaaacca 1080
gtctgctcgc tgccctgttt ccgccaccga cccattcccg atgcgacgtc acccgacttc 1140
gcccttcgt agcgcctccg tcccttgctc tgtttttctc gtgacgtcga tccgctcgac 1200
aaattagatc tcgctcgtcg caacgatatg acttatcccc tgcggtcggg ggacttgacg 1260
aactagaaaa ttatggctct catgtggtct ttttgcccc tcctaaattt tcgtcctgcg 1320
tccgccactc tatagctcgc ggagtgcggg gtaccgtatc gtgaatctga agtcgaacag 1380
tgatggcgta ctatctaata tcccgctgag taatatcact gttgccgacg atgggaatct 1440
ctagttttga cagaaaccaa ggcaactgct atgctagcta attaatcca gagagatcga 1500
tttctacagt gctagctgca acaatcgatg caattggcat cagacgatat atgctaattg 1560
ttttctttat cgatacgtgg tcaacagtgc gatccagatt acagagttct ctgcacctga 1620
tcggatctca tcgcacatgg acacccatct gccaacccaa cgcggggcg ggaacccgaa 1680

```

09/786835 061901

acatcgcgtc catgcacgac cccacgcag ctataaatac ccatgcaatg caatgcagcg 1740
ggtcatcatc atcgactcca 1760

<210> 2
<211> 1348
<212> DNA
<213> Zea mays

<400> 2
ttgggattgg gtttgtgcgg acgcccactg gttcttggct ctgctggagt cttgtctgtg 60
gagctggttg tgcgctgct aacaatctct tgttggttgc gacgacttgc cgtggccacg 120
cgttaactgc gctctgctca aagaaattgc ctatacagtg cctggcgtca gctgcagcg 180
ttgaatccga ggtcgccagt acgaggcaaa cgctgagctg catgctgtac cggctttatt 240
aataactgag ctctactcgc tacgttagct ataggcacca gaggcggatc cagagggggg 300
gcaaagtagg ccatggcccc cctcaatttt tgtacaccct tttataccta atgattattt 360
agtattaagt gtagaaaaaa tcaaaatttt agatagtata aaaatcttgt tctaaatctc 420
taaaactaat agctagttgc taaaatttag caaaagggtt agaatggatc aactaattgt 480
tagctatacc ctttaaataa gtattaattg ttagatggcc caacctagct aaaaccagtt 540
aacagtttag tctagatggt tagaacatgg cctaaacaac catattatga gcctactcgc 600
cggatcctct cggttcagtc ctagecatct ccagctctgac tctaactctc ccgacaccac 660
ctaaaccagt ctgctcgcgt cctgttttcc gccaccgacc cattcccgat gcgacgtcac 720
ccgacttcgc ccttcgttag cgctccgctc ccttgtctct tttttctcgt gacgtcgatc 780
cgctcgacaa attagatctc gctcgtcgca acgatatgac ttatcccctg cggtcgggtg 840
acttgacgaa ctagaaaatt atggctctca tgtggtcttt ttggccctc ctaaaatttc 900
gtcctgcgtc cgccactcta tagctgcggg agtgcgggg accgtatcgt gaactctgaag 960
tcgaacagtg atggcgtact atctaattgc ccgtgcagta atatcactgt tgccgacgat 1020
gggaatctct agttttgaca gaaaccaagg caactgctat gctagctaata taattccaga 1080
gagatcgatt tctacagtgc tagctgcaac aatcgatgca attggcatca gacgatatat 1140
gctaattggt ttctttatcg atacgtggtc aacagtgcga tccagattac agagttctct 1200
cgccctgacg ggatctcatc gcacatggac acccatctgc caaccacaac cggggcgggg 1260
aaccggaaac atcgcgcca tgcacgaccc ccacgcagct ataaataccc atgcaatgca 1320
atgcagcggg tcatcatcat cgactcca 1348

<210> 3
<211> 1096
<212> DNA
<213> Zea mays

<400> 3
ctactcgcta cgttagctat aggcaccaga ggccgatcca gagggggggc aaagtagggc 60
atggcccccc ctcaattttg tacacccttt tatacctaata gattatttag tattaagtgt 120
agaaaaaatac aaaatttttag atagtataaa aatcttggtc taaatctcta aaactaatag 180
ctagttgcta aaatttagcca aaaggtttag aatggatcaa ctaattgtta gctataccct 240
ttaaataagt attaattggt agatggccca acctagctaa aaccagttaa cagttagctc 300
tagatgttta gaacatggcc taaacaacca tattatgagc ctactcgccg gatcctctcg 360
gttcagtcct agcatctacc agtctgactc taatcttccc gacaccacct aaaccagtct 420
gctcgtcgcc ctgtttccgc caccgacca ttcccgatgc gacgtcaccc gacttcgccc 480
cttcgtagcg cctcgtccc ttgtcctgtt tttctcgtga cgtcgatccg ctcgacaaat 540
tagatctcgc tcgtcgcaac gatatgactt atccctcgcg gtcgggtggac ttgacgaact 600
agaaaaattat ggctctcatg tggctttttt ggccctcctt aaattttcgt cctgcgtccg 660
ccactctata gctgcgggag tgcggggtac cgtatcgtga atctgaagtc gaacagtgat 720

```

ggcgactat ctaatgtccc gtgcagtaat atcactgttg cgcacgatgg gaatctctag 780
ttttgacaga aaccaaggca actgctatgc tagctaatta attccagaga gatcgatttc 840
tacagtgcta gctgcaacaa tcgatgcaat tggcatcaga cgatatatgc taatgggtttt 900
ctttatcgat acgtgggtcaa cagtgcgata cagattacag agttctctcg cctgatcgcg 960
atctcatcgc acatggacac ccatctgcc aaccaacgcg gggcggggaa cccgaaacat 1020
cgcgtccatg cagcaccccc acgcagctat aaatacccat gcaatgcaat gcagcgggtc 1080
atcatcatcg actcca 1096

```

<210> 4
 <211> 746
 <212> DNA
 <213> Zea mays

```

<400> 4
gatcctctcg gttcagtcct agcatctacc agtctgactc taatcttccc gacaccacct 60
aaaccagtct gctcgtgccc ctgtttccgc caccgaacca ttcccgatgc gacgtcacc 120
gaattcgccc cttcgtagcg cctccgtccc ttgtcctggt tttctcgtga cgtcgatccg 180
ctcgacaaat tagatctcgc tcgtcgcaac gatatgactt atcccctgcg gtcgggtggac 240
ttgacgaact agaaaattat ggctctcatg tggctctttt ggcccctcct aaattttcgt 300
cctgcgtccg ccaactctata gctgcgggag tgcggggtac cgtatcgtga atctgaagtc 360
gaacagtgat ggcgactat ctaatgtccc gtgcagtaat atcactgttg cgcacgatgg 420
gaatctctag ttttgacaga aaccaaggca actgctatgc tagctaatta attccagaga 480
gatcgatttc tacagtgcta gctgcaacaa tcgatgcaat tggcatcaga cgatatatgc 540
taatgggttt ctttatcgat acgtgggtcaa cagtgcgata cagattacag agttctctcg 600
ccctgatcgg atctcatcgc acatggacac ccatctgcc aaccaacgcg gggcggggaa 660
cccgaacat cgcgtccatg cagcaccccc acgcagctat aaatacccat gcaatgcaat 720
gcagcgggtc atcatcatcg actcca 746

```

<210> 5
 <211> 554
 <212> DNA
 <213> Zea mays

```

<400> 5
gatctcgtc gtcgcaacga tatgaattat cccctgcggt cgggtggactt gacgaactag 60
aaaattatgg ctctcatgtg gtcttttttg cccctcctaa attttcgtcc tgcgtccgcc 120
actctatagc tgcgggagtg cgggggtacc tatcgtgaat ctgaagtcga acagtgatgg 180
cgtactatct aatgtcccgt gcagtaatat cactgttgcc gacgatggga atctctagtt 240
ttgacagaaa ccaaggcaac tgctatgcta gctaattaat tccagagaga tcgatttcta 300
cagtgttagc tgcaacaatc gatgcaattg gcatcagacg atatatgcta atgggtttct 360
ttatcgatac gtggtcaaca gtgcgatcca gattacagag ttctctcgcc ctgatcggat 420
ctcatcgac atggacaccc atctgccaac ccaacgcggg gcggggaacc cgaaacatcg 480
cgtccatgca cgacccccac gcagctataa ataccatgc aatgcaatgc agcgggtcat 540
catcatcgac tcca 554

```

<210> 6
 <211> 406
 <212> DNA
 <213> Zea mays

098633-06191

<400> 6

```
cgtatcgtga atctgaagtc gaacagtgat ggcgtactat ctaatgtccc gtgcagtaat 60
atcactgttg ccgacgatgg gaatctctag ttttgacaga aaccaaggca actgctatgc 120
tagctaatta attccagaga gatcgatttc tacagtgccta gctgcaacaa tcgatgcaat 180
tggcatcaga cgatatatgc taatggtttt ctttatcgat acgtgggtcaa cagtgcgcatc 240
cagattacag agttctctcg cccatgatcg atctcatcgc acatggacac ccatctgcca 300
acccaacgcg gggcggggaa cccgaaacat cgcgtccatg cagaccccc acgcagctat 360
aatatcccat gcaatgcaat gcagcgggtc atcatcatcg actcca 406
```

<210> 7

<211> 190

<212> DNA

<213> Zea mays

<400> 7

```
cgatacgtgg tcaacagtcg gatccagatt acagagttct ctgcacctga tcggatctca 60
tcgcacatgg acacccatct gccaacccaa cgcggggcgg ggaacccgaa acatcgcgtc 120
catgcacgac cccacgcag ctataaatac ccatgcaatg caatgcagcg ggtcatcatc 180
atcgactcca 190
```

<210> 8

<211> 1354

<212> DNA

<213> Zea mays

<400> 8

```
ctcgagatgt tggatagaag aggaattgcg agcaaccgag gtctccaagg caccgcaacg 60
cgcacacctg gcagtgaggt ggtcgctgat gtcgcgaagg agcttcgatt gttcgctgat 120
cgcgagcttc agatttggtt ccatgacccc agattgtttg gcgcggcgcg tgtaatgagc 180
agacgccaga cccgagaccg gtgtctccga taccaattgt tagcaatagc agcggacaag 240
cagcgaacaa tccgagagac atagtgtggg ggaggagcag atagcttgga gaggaagaag 300
aacaatagtg acagcgtaga ttctctctcg gtctctctcg ttacctctc tcaatctaata 360
actcctatta caaggtcggc ccatcaggcg acccagtcgc ggttcacaga acttgggatt 420
gggttttgtg ggacgccgac tggttcttgg ctctgctgga gtcttgtctg tggagctggt 480
gctcgcgctg ctaacaatct cttgttgttt gcgacgactt cgcgtggcca cgcgttaact 540
gogctctgct caaagaaatt gcctatacag tgcctggcgt cagctgcagg cgttgaatcc 600
gaggtcgcca gtacgaggca aacgctgagc tgcctgctgt accggcttta ttaataactg 660
agctctactc gctacgttag ctataggcac cagaggcgga tccagagggg gggcaaagta 720
ggtatagaaa aatcaaaatt ttagatagta taaaaatctt gttctaaatc tctaaaacta 840
atagctagtt gctaaaatta gccaaaaggt ttagaatgga tcaactaatt gttagctata 900
ccctttaaat aagtattaat tgtagatgg cccaacctag ctaaaaccag ttaacagtta 960
gctctagatg tttagaacat ggcctaaaca accatattat gagcctactc gccggatcct 1020
ctcggttcag tcttagcatc taccagtctg actctaactc tcccgacacc acctaaacca 1080
gtctgctcgc tgccctgttt ccgccaccga cccattcccg atgcgacgtc acccgacttc 1140
gccccttctg agcgcctcgc tcccttgtcc tgtttttctc gtgacgtcga tccgctcgac 1200
aaattagatc tcgctcgctc caacgatatg acttatcccc tgcggtcggg ggacttgacg 1260
aactagaaaa ttatggctct catgtggtct ttttgcccc tcctaaaatt tcgtcctcgc 1320
tccgccactc tatagctgcg ggagtgcggg gtac 1354
```

<210> 9
 <211> 1497
 <212> DNA
 <213> Zea mays

<400> 9
 ctcgagatgt tggatagaag aggaattgag agcaaccgag gtctccaagg cacgccaacg 60
 cgcacccctgc gcagttaggt ggtcgctgat gtcgcgaagg agcttcgatt gttcgtcgat 120
 cgcgagcttc agatttggct ccatgacccc agattgtttg gcgcggcgcg tgtaatgagc 180
 agacgccaga cccgagaccg gtgtctccga taccatttgt tagcaatagc agcggacaag 240
 cagcgaacaa tccgagagac atagtttggg ggaggagcag atagcttgga gaggaagaag 300
 aacaatagtg acagcgtaga ttctctctctg gtctctctctg ttacctctctc tcaatctaata 360
 actcctatta caaggctcggc ccacagggcg acccagtcgc ggttccacga acttgggatt 420
 gggtttgtgc ggacgccgac tggttcttgg ctctgctgga gtcttgtctg tggagctgtt 480
 gctcgcgctg ctaacaatct cttgttgttt gcgacgactt cgcgtggcca cgcgttaact 540
 gcgctctgct caaagaaatt gcctatacag tgcctggcgt cagctgcagg cgttgaatcc 600
 gaggtcgcca gtacgaggca aacgctgagc tgcctgctgt accggcttta ttaataactg 660
 agctctactc gctacgttag ctataggcac cagaggcgga tccagagggg gggcaaagta 720
 ggccatggcc cccctcaat tttgtacacc cttttatacc taatgattat ttagtattaa 780
 gtgtagaaaa aatcaaaatt ttagatagta taaaaatctt gttctaaatc tctaaaacta 840
 atagctagtt gctaaaatta gccaaaaggt ttagaatgga tcaactaatt gtttagctata 900
 ccctttaaat aagtattaat tgttagatgg cccaacctag ctaaaaccag ttaacagtta 960
 gctctagatg tttagaacat ggcctaaaca accatattat gagcctactc gccggatcct 1020
 ctcggttcag tcttagcctc taccagctctg actctaactc tcccgacacc acctaaacca 1080
 gtctgctcgc tgccctgttt ccgccaccga cccattcccg atgcgacgct acccgacttc 1140
 gccctctcgt agcgcctccg tcccttctcc gctttttctc gtgacgctga tccgctcgac 1200
 aaattagatg tcgctcgtcg caacgatatg acttatcccc tgcggtcggg ggacttgacg 1260
 aactagaaaa ttatggctct catgtggtct ttttggtccc tcttaaattt tgcctctcgc 1320
 tccgccactc tatagctcgc ggagtgcggg gtaccgtatc gtgaatctga agtcgaacag 1380
 tgatggcgta ctatctaata tcccgctcag taatatcact gttgccgacg atgggaatct 1440
 ctagttttga cagaaaccaa ggcaactgct atgctagcta attaatcca gagagat 1497

<210> 10
 <211> 1550
 <212> DNA
 <213> Zea mays

<400> 10
 ctcgagatgt tggatagaag aggaattgag agcaaccgag gtctccaagg cacgccaacg 60
 cgcacccctgc gcagttaggt ggtcgctgat gtcgcgaagg agcttcgatt gttcgtcgat 120
 cgcgagcttc agatttggct ccatgacccc agattgtttg gcgcggcgcg tgtaatgagc 180
 agacgccaga cccgagaccg gtgtctccga taccatttgt tagcaatagc agcggacaag 240
 cagcgaacaa tccgagagac atagtttggg ggaggagcag atagcttgga gaggaagaag 300
 aacaatagtg acagcgtaga ttctctctctg gtctctctctg ttacctctctc tcaatctaata 360
 actcctatta caaggctcggc ccacagggcg acccagtcgc ggttccacga acttgggatt 420
 gggtttgtgc ggacgccgac tggttcttgg ctctgctgga gtcttgtctg tggagctgtt 480
 gctcgcgctg ctaacaatct cttgttgttt gcgacgactt cgcgtggcca cgcgttaact 540
 gcgctctgct caaagaaatt gcctatacag tgcctggcgt cagctgcagg cgttgaatcc 600
 gaggtcgcca gtacgaggca aacgctgagc tgcctgctgt accggcttta ttaataactg 660
 agctctactc gctacgttag ctataggcac cagaggcgga tccagagggg gggcaaagta 720
 ggccatggcc cccctcaat tttgtacacc cttttatacc taatgattat ttagtattaa 780
 gtgtagaaaa aatcaaaatt ttagatagta taaaaatctt gttctaaatc tctaaaacta 840

0986335 061901

```

atagctagtt gctaaaatta gccaaaaggt ttagaatgga tcaactaatt gttagctata 900
ccctttaaat aagtattaat tgttagatgg cccaacctag ctaaaaccag ttaacagtta 960
gctctagatg tttagaacat ggccataaaca accatattat gagcctactc gccggatcct 1020
ctcggttcag tcctagcatc taccagtctg actctaactc tcccgcacacc acctaaacca 1080
gtctgctcgc tgccctgttt ccgccaccga cccattcccg atgcgacgtc acccgacttc 1140
gccccttcgt agcgccctccg tcccttgtec tgttttttctc gtgacgtcga tccgctcgac 1200
aaattagatc tcgctcgctc caacgatatg acttatcccc tgcggtcggt ggacttgacg 1260
aactagaaaa ttatggctct catgtggtct ttttggcccc tcctaaattt tcgtcctgcg 1320
tccgccactc tatagctgcg ggagtgcggg gtaccgtatc gtgaatctga agtcgaacag 1380
tgatggcgta ctatctaata tcccgtcgag taatatcact gttgccgacg atgggaatct 1440
ctagttttga cagaaaccaa ggcaactgct atgctagcta attaattcca gagagatcga 1500
tttctacagt gctagctgca acaatcgatg caattggcat cagacgatat 1550

```

<210> 11
 <211> 1710
 <212> DNA
 <213> Zea mays

```

<400> 11
ctcgagatgt tggatagaag aggaattgcg agcaaccgag gtctccaagg cagccaacg 60
cgcacccctgc gcagttaggt ggtcgctgat gtcgcgaagg agcttcgatt gttcgtcgat 120
cgcgagcttc agatttggct ccatgacccc agattgtttg gcgcggcgcg tgtaatgagc 180
agacgccaga cccgagaccg gtgtctccga taccaattgt tagcaatagc agcggacaag 240
cagcgaacaa tccgagagac atagtttggg ggaggagcag atagcttgga gaggaagaag 300
aacaatagtg acagcgtaga ttctctctct gtctcctctg ttacctctc tcaatctaact 360
actcctatta caaggtcggc ccatcaggcg acccagtcgc ggttccacga acttgggatt 420
gggtttgtgc ggacgccgac tggttcttgg ctctgctgga gtcttgtctg tggagctgtt 480
gctcgcgctg ctaacaatct cttgttgttt gcgacgactt cgcgtggcca cgcgttaact 540
gcgctctgct caaagaaatt gcctatacag tgcctggcgt cagctgcagg cgttgaatcc 600
gaggctgcca gtacgaggca aacgctgagc tgcctgctgt accggcttta ttaataactg 660
agctctactc gctacgttag ctataggcac cagaggcgga tccagagggg gggcaaagta 720
ggccatggcc cccctcaat tttgtacacc cttttatacc taatgattat ttagtattaa 780
gtgtagaaaa aatcaaaatt ttagatagta taaaaatctt gttctaaato tctaaaacta 840
atagctagtt gctaaaatta gccaaaaggt ttagaatgga tcaactaatt gttagctata 900
ccctttaaat aagtattaat tgttagatgg cccaacctag ctaaaaccag ttaacagtta 960
gctctagatg tttagaacat ggccataaaca accatattat gagcctactc gccggatcct 1020
ctcggttcag tcctagcatc taccagtctg actctaactc tcccgcacacc acctaaacca 1080
gtctgctcgc tgccctgttt ccgccaccga cccattcccg atgcgacgtc acccgacttc 1140
gccccttcgt agcgccctccg tcccttgctc tgttttttctc gtgacgtcga tccgctcgac 1200
aaattagatc tcgctcgctc caacgatatg acttatcccc tgcggtcggt ggacttgacg 1260
aactagaaaa ttatggctct catgtggtct ttttggcccc tcctaaattt tcgtcctgcg 1320
tccgccactc tatagctgcg ggagtgcggg gtaccgtatc gtgaatctga agtcgaacag 1380
tgatggcgta ctatctaata tcccgtcgag taatatcact gttgccgacg atgggaatct 1440
ctagttttga cagaaaccaa ggcaactgct atgctagcta attaattcca gagagatcga 1500
tttctacagt gctagctgca acaatcgatg caattggcat cagacgatat atgctaattg 1560
ttttctttat cgatacgtgg tcaacagtgc gatccagatt acagagttct ctgccctga 1620
tcggatctca tcgcacatgg acacccatct gccaaaccaa cgcggggcgg ggaacccgaa 1680
acatcgctgc catgcacgac cccacgcgac 1710

```

<210> 12
 <211> 856

09786835 061501

<212> DNA

<213> Zea mays

<400> 12

```

gtttccctgc ggcattgcagc tggagcgctc ggcgttagtg tcagtccatt ttggagtcgg 60
ttcgtgtacc tgagttttcc ttgaggagat gctgggtttc gctttctttg ttgcaagaat 120
tattagtcag gaataatggg cctcgcgtga ttgtagtctg agtaggtagg tagtccagca 180
tgcaactgac atgtaatggc tactacgagt atatgcttgc atcgccatct tttatttgct 240
gtgccccctg ttgttacagt gttcctcatc gatctttaga ggtcgttgct accgggtttt 300
atcttccaaa acacaaaata tactggaacg actcttgagc tctcttgcta ctgttgaatg 360
tgactattga ctagctaatt agctatagta gcatttatac agttgttgct aattattata 420
tatagggttg gaaaaaaaaa ctcgagctcg acgacttaka ttgggtctcg cagcagctcc 480
agctcggctt gactcggact ggctcgacgc tcggacgagc ccgagccgag cctatttttg 540
tggctcgtga aaagagcgag ccaactcggc tcgctccaac tcgcgagcta gctcgtggct 600
cgaggttaaca acaattttta catataattc taattagcat ataattattg taccgagtat 660
acaattaatt tatgttattt gagattacta tacaatttaa tctagtatta gtgatatatc 720
tattttacta atttaaaata tgttatttaa aatattttta agattttata ttataattta 780
gagagcagat ctacgtttat agctagtacc gagctgcttt ttccagctca ccagatgggt 840
gcgccgaacc gagctc

```

<210> 13

<211> 440

<212> DNA

<213> Zea mays

<400> 13

```

gtttccctgc ggcattgcagc tggagcgctc ggcgttagtg tcagtccatt ttggagtcgg 60
ttcgtgtacc tgagttttcc ttgaggagat gctgggtttc gctttctttg ttgcaagaat 120
tattagtcag gaataatggg cctcgcgtga ttgtagtctg agtaggtagg tagtccagca 180
tgcaactgac atgtaatggc tactacgagt atatgcttgc atcgccatct tttatttgct 240
gtgccccctg ttgttacagt gttcctcatc gatctttaga ggtcgttgct accgggtttt 300
atcttccaaa acacaaaata tactggaacg actcttgagc tctcttgcta ctgttgaatg 360
tgactattga ctagctaatt agctatagta gcatttatac agttgttgct aattattata 420
tatagggttg gaaaaaaaaa

```

<210> 14

<211> 3416

<212> DNA

<213> Zea mays

<400> 14

```

ctcgagatgt tggatagaag aggaattgct agcaaccgag gtctccaagg cagccaacg 60
cgcactctgc gcagttaggt ggtcgtgat gtcgcgaagg agcttcgatt gttcgtcgat 120
cgcgagcttc agatttggtt ccatgacccc agattgtttg gcgcggcgcg tgtaatgagc 180
agacgccaga cccgagaccg gtgtctccga taccaattgt tagcaatagc agcggacaag 240
cagcgaacaa tccgagagac atagtttggt ggaggagcag atagcttgga gaggaagaag 300
aacaatagtg acagcgtaga ttctctctct gtctctctct ttaccctctc tcaatctaatt 360
actcctatta caaggctggc ccatcaggcg acccagtcgc ggttccacga acttgggatt 420
gggtttgtgc ggacgcgcag tggttcttgg ctctgctgga gtcttgctct tggagctggt 480
gctcgcgctg ctaacaatct cttgttgttt gcgacgactt cgcgtggcca cgcgttaact 540
gcgctctgct caaagaaatt gcctatacag tgcttgccgt cagctgcagg cgttgaatcc 600

```


gaggtcgcca	gtacgaggca	aacgctgagc	tgcattgctgt	accggcttta	ttaataactg	660
agctctactc	gctacgttag	ctataggcac	cagaggcgga	tccagagggg	gggcaaagta	720
ggccatggcc	ccccctcaat	tttgtacacc	ctttttatacc	taatgattat	ttagtatttaa	780
gtgtagaaaa	aatcaaaaatt	ttagatagta	taaaaatctt	gttctaaatc	tctaaaacta	840
atagctagtt	gctaaaaatta	gccaaaagg	ttagaatgga	tcaactaatt	gttagctata	900
ccctttaaat	aagtattaat	tgtagatgg	cccaacctag	ctaaaaccag	ttaacagtta	960
gctctagatg	tttagaacat	ggcctaaaca	accatattat	gagcctactc	gccggatcct	1020
ctcgggtcag	tcctagcatc	taccagtctg	actctaattc	tcccgacacc	acctaaacca	1080
gtctgctcgc	tgccctgttt	ccgccacoga	cccattcccg	atgcgacgtc	acccgacttc	1140
gccccttcgt	agcgctctgt	tcccttgctc	tgtttttctc	gtgacgtcga	tccgctcgac	1200
aaattagatc	tcgctcgctg	caacgatatg	acttatcccc	tgcggtcggg	ggacttgacg	1260
aactagaaaa	ttatggctct	catgtggctc	ttttggcccc	tcctaaattt	tcgtcctgcg	1320
tccgccactc	tatagctgcg	ggagtgcggg	gtaccgtatc	gtgaatctga	agtcgaacag	1380
tgatggcgta	ctatctaatt	tcccgtcgag	taatatcact	gttgccgacg	atgggaatct	1440
ctagttttga	cagaaaccaa	ggcaactgct	atgctagcta	attaattcca	gagagatcga	1500
tttctacagt	gctagctgca	acaatcgatg	caattggcat	cagacgatat	atgctaattg	1560
ttttctttat	cgatacgtgg	tcaacagtgc	gatccagatt	acagagttct	ctcgccctga	1620
tcggatctca	tcgcacatgg	acacccatct	gccaaaccaa	cgcgggggcg	ggaacccgaa	1680
acatcgcgtc	catgcaacgac	ccccacgcag	ctataaatac	ccatgcaatg	caatgcagcg	1740
ggtcatcatc	atcgactcca	cctggactcg	ctcactggca	atggctacca	gcaccaagct	1800
cgtggccctt	ggcttgcccg	tccctgctgag	cgttggttcc	tccgatgcag	cgagggtagc	1860
taggcttggg	agctatgcgt	ctgccggagg	tggcgggcgg	ggtggaggcg	gcggctcccg	1920
atcgaccggg	gctgggtacg	gcgggtgggtc	tggcggtggc	ggtggctacg	gcataggcaa	1980
gggtggcggg	gactggtgga	acaatttcgt	ctccagcgta	gcggggcggtg	gtggtggggg	2040
gcaaggcgga	ggaggtggca	ccaacgggtg	atccgggtcc	ggcggtgggt	cgggctatgg	2100
ttctggcacc	agttccaccc	cagcctctgg	ccccagcagt	ggcaactacg	ccaacgctga	2160
gggcaagggc	gcgggtgggg	gcatggggcg	tgggtgcgac	gcccgaacgc	cctacggttc	2220
tgggtgccgg	ggaggcggtt	gcaaggggcca	gggcgagagt	ggcgctcgat	tggcacgcgc	2280
gtccgatggc	tactacaatg	gcggcgccgc	tgatgctacc	ggtggtgggt	ccggcgctgg	2340
cgggtggacat	ggcggtggcg	cagccggagc	tccaagctac	gggactggag	gtggccttgg	2400
cggaggcaag	ggcgaggccg	gcagcgacgg	gtcctggggg	tcaggttacg	cggcgggaat	2460
cgggtgccggc	actggcggcg	gcgggtggcg	cggatcgcaa	ggcggtctct	gtggtggcgg	2520
cggctctggg	tccggtatcc	gcagcggtgg	aatccactga	gtttccctgc	ggcatgcagc	2580
tggagcgctc	ggcgtagtg	tcagtcocatt	ttggagtcgg	ttcgtgtacc	tgagttttcc	2640
ttgaggagat	gctggttttc	gctttctttg	ttgcaagaat	tattagtcag	gaataatggg	2700
cctcgcgctga	ttgtagttcg	agtaggtagg	tagtccagca	tgcaactgac	atgtaatggc	2760
tactacgagt	atatgcttgc	atcgccatct	tttatttgct	gtgccccttg	ttgttacagt	2820
gttctctatc	gatctttaga	ggtcgttgct	accggttttt	atcttccaaa	acacaaaata	2880
tactggaacg	actcttgacg	totcttgcta	ctggtgaatg	tgactattga	ctagctaatt	2940
agctatagta	gcatttatac	agttgttgct	aattattata	tatagggttg	gaaaaaaaaa	3000
ctcgagctcg	acgacttaga	ttgggtctcg	cagcagctcc	agctcggtct	gactcgact	3060
ggctcgacgc	tcggacgagc	ccgagccgag	cctatttttg	tggctcgtga	aaagagcgag	3120
ccaactcggc	tcgctccaac	tcgcgagcta	gctcgtggct	cgaggtaaca	acaattttta	3180
catataattc	taattagcat	ataatattag	taccgagtat	acaattaatt	tatgttattt	3240
gagattacta	tacaaaattaa	tctagtatta	gtgatataat	tattttacta	attttaaaata	3300
tgttatttta	aatattttta	agattttata	ttataattta	gagagcagat	ctacgtttat	3360
agctagtacc	gagctgcttt	ttccagctca	ccagatgggt	gcgccgaacc	gagctc	3416

<210> 15
 <211> 1074
 <212> DNA
 <213> Zea mays

09785935.061901

ggctcggtgc	atgagcgagc	cgagtcgaagc	tggggttggc	tcgtttccac	ccctagttat	60
atacgatcaa	attggaccgc	acgacactga	cattaacact	attaaacata	acactattat	120
taatcggtgc	gtgctatatt	agcattagtgc	actagctatc	ggtacagaca	tgacactata	180
ttgcctaata	atgtcggtgc	tgacactata	tagaggggtat	ctctagtagt	ttttctatct	240
tagttcctat	ctgtcggtac	cctgaaacta	gggtatccct	tactattgta	taaagacgca	300
gtacccacgc	ggctatcttt	agacgcgtgc	taaacgagct	gtatgtggga	ccaggccatg	360
actcgcccca	gcctcggggc	actactctag	gccagcaaca	gcgcttgacc	ccaccacatg	420
ggcgggtccg	cggccgccac	gtgtccagag	aaagtgatata	actccaaggc	atcaacagtg	480
agtcgggaca	tgggagagtg	tgggacccat	gcgggacccc	tgtatatacg	gtccaggcct	540
ccaagtttgg	cccaggacct	taggccattg	ccgctgtgcg	ccggcgggtg	aatgagtatc	600
atgtcgtagc	tgccgtcgag	ggacatgaac	tcaagactcc	cgaacaggag	caccttcccg	660
ggttgagagc	gttgctcgag	actgcccatc	tggagcttga	cgggaagctg	ttcgtcaaca	720
cgcagcaggc	ccctactgtg	cgcgccaaact	gtcggcgctt	cgagaccggg	gggtccctaa	780
gccgacgagt	gaatctcgcc	gcgtgcccc	gccagatgg	gtcgacgcga	ggccgagcgc	840
gaagggggga	agtgaggtgc	ccggaagatag	gcgtgagaga	gggtcaaatc	ccgcggcctc	900
cgtgttcgtc	tcgcgcccag	gtcgggtgcg	cttgcagtag	ggggttacaa	gcgtccacgc	960
gggagaggga	gcaagcggcc	taacacgagc	gcattgtctc	tcctcgtccc	cgcgcggcca	1020
accctatcta	agagggccct	ggtccttctc	tttataggcg	taaggagagg	atcc	1074